

A version of Goodwillie calculus for non-cubes

Robin Stoll

Born October 3, 1996 in Bonn, Germany

September 27, 2019

Master's Thesis Mathematics

Advisor: Dr. Emanuele Dotto

Second Advisor: Prof. Dr. Stefan Schwede

MATHEMATICAL INSTITUTE

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN

Contents

Introduction	2
1. Notation and conventions	6
2. Excisive functors	11
3. Maps between approximations	28
3.1. Indirect maps	28
3.2. Direct maps	30
3.3. Example application: punctured cubes	31
4. The structure of shapes	34
4.1. Full shapes	34
4.2. Free shapes	36
4.3. Boring shapes	39
5. The Taylor graph	45
5.1. Relation to the Taylor tower	50
Appendices	52
A. The calculus of mates	52
B. Basic ∞-categorical facts	55
C. Tools for Kan extensions and (co)limits	56
C.1. Kan extensions	56
C.2. (Co)Limits	59
C.3. Preservation of Kan extensions and (co)limits	63
D. Generalities	69
D.1. about posets	69
D.2. about ∞ -categories	70
D.3. about cartesian diagrams	72

Introduction


Classical Goodwillie calculus was developed by Tom Goodwillie in the series of papers [Goo90; Goo92; Goo03] as a tool to study functors from spaces to spaces or spectra. Throughout the subsequent years it turned out to be a very fruitful theory that has, together with a few variations, found many applications in homotopy theory and elsewhere, e.g. to algebraic K-theory, chromatic homotopy theory, or embedding spaces of manifolds. A survey of some of these can be found in [AC]. Moreover there are generalizations to the setting of model categories (see e.g. [Kuh] or [Per]) and to the setting of quasi-categories (see [LurHA, Section 6]). The latter framework is the one we will use in this thesis.

The notion most fundamental to the theory is the one of an n -excisive functor, that is one that sends cubical diagrams that are *strongly cocartesian* to ones that are *cartesian*, i.e. a homotopy limit diagram. Strongly cocartesian is a slightly more complicated condition: it means that every 2-dimensional face of the cube is cocartesian, i.e. a homotopy pushout. The importance of this notion lies in the fact that any functor F can be approximated by functors that are n -excisive. This is done via an explicit construction called the *Taylor tower* of F which is a tower

$$F \longrightarrow \dots \longrightarrow P_2 F \longrightarrow P_1 F \longrightarrow P_0 F$$

lying under F built from universal n -excisive approximations to F , denoted $P_n F$, and maps between them induced by their universal properties. In good cases this tower *converges*, i.e. reconstructs F in the sense that the canonical map $F \rightarrow \text{holim}_n P_n F$ is an equivalence. Moreover, it possesses a wealth of structure that can be used to analyze the functor F , even in the most basic case when F is the identity. A short overview over the current state of art in this direction can again be found in [AC].

Now, the starting point of this thesis is an equivalent definition of a strongly cocartesian cubical diagram: one can instead require the diagram to be a left Kan extension of its restriction to the initial star of the cube, i.e. along an inclusion as in the following picture (for the case of a 3-cube):



This reformulated condition does not make any reference to the structure of the cube; it only needs the datum of the above inclusion. In particular this allows us to generalize the notion to arbitrary maps of posets $s: \mathcal{P} \rightarrow \mathcal{Q}$ by defining a \mathcal{Q} -indexed diagram to be *s-cocartesian* if it is a left Kan extension along s of its restriction along s . If \mathcal{Q} additionally has an initial object, then such a condition induces a notion of excision:

Definition. A functor is *s-exciseive* if it sends *s*-cocartesian diagrams to cartesian diagrams.

The goal of this thesis is to study this notion, in particular in the context of Goodwillie calculus.

One of the first questions to ask is whether the construction of the universal excisive approximation still works. Indeed, our first main result states that this is the case for a class of well-behaved maps of posets, called *shapes* (see Definition 2.34), which generalize the inclusions of the classical setting:

Theorem. *Let σ be a full shape and F a functor. Then there is an explicit construction of a functor $P_\sigma F$ such that it is a universal σ -excisive approximation to F .*

(See Theorem 2.42 for the precise statement.) Here a shape is *full* if it is full as a functor.

Clearly one would also like to have an analogue of the Taylor tower in this more general framework. Again, this exists (though it is a bit more technical to construct) and takes the form of the *Taylor graph*: a diagram that contains, for a fixed functor F , all of its universal excisive approximations $P_\sigma F$ and the maps between them induced by their universal properties. Large parts of the thesis are devoted to studying this diagram. The main result we will prove is the following:

Theorem. *The limits of the Taylor tower and the Taylor graph always agree when the latter is restricted to non-boring shapes between finite posets.*

(Actually we will even show the stronger statement that an evident functor between their indexing categories is homotopy initial, see Theorem 5.18.) Notably this tells us that any convergence criteria for the Taylor tower can also be used for the Taylor graph. Here, a *boring* shape is one that fulfills a certain combinatorial condition. They are so called because it turns out that, in the finite case, any functor is excisive with respect to them. In particular we do not lose any information by discarding them.

In proving this theorem we first pursue the naturally arising question of when there exists a map $P_\sigma F \rightarrow P_\tau F$ in the Taylor graph or, equivalently, of when τ -excisive implies σ -excisive. It turns out that it is not true that any (naively defined) morphism of shapes $f: \tau \rightarrow \sigma$ induces such a map. However we are able to give an explicit combinatorial condition on f for this to be the case. Moreover, maybe surprisingly, it turns out that there is a second, independent condition one can put on f that guarantees the existence of a map $P_\tau F \rightarrow P_\sigma F$, i.e. this time in the other direction. Together, these two conditions turn out to be very useful in studying shapes and their relations.

Further pursuing a proof of the second main theorem, we are lead to the question of when, for a shape σ , there exists an n such that σ -excisive is equivalent to n -excisive. Using the tools outlined in the previous paragraph we are able to give a partial answer: if σ is a *free* shape generated by a finite poset, then there is such an n . This is particularly useful since it turns out that excision with respect to a non-boring full shape implies excision for a free shape. Together with showing that, in the finite case, one can reduce to the case of full shapes, this is enough to prove the theorem.

We finish by posing the (probably) most important question left unanswered: is the inclusion of the Taylor tower into the Taylor graph an equivalence? On the answer depends the usefulness of the framework. If it were yes, the graph would not contain more information than just the tower and we would have found evidence that the cubes are the correct indexing diagrams to study. If it were no, we would have obtained a finer resolution of the tower that could potentially contain more useful information.

The following is a short overview over the distribution of the contents over the sections and appendices. In Section 1 we collect the conventions and notations we will use throughout the rest of the thesis. In Section 2 we define f -excisive and shapes, give the construction of the universal excisive approximation, and prove the first main theorem. In Section 3 we give the conditions for morphisms of shapes to induce maps between their universal excisive approximations. In Section 4 we employ these conditions to study shapes and their relations, introducing the notions of free and boring shapes along the way. In Section 5 we construct the Taylor graph and use the results of the previous two sections to prove the second main theorem. In Appendix A we recall the calculus of mates of natural transformations. In Appendix B we recall a number of ∞ -categorical facts that will be used throughout the thesis (often without reference). In Appendix C we give references or proofs for these and other needed basic facts about (co)limits and Kan extensions. In Appendix D we prove various general facts that are needed but would hinder the flow of the main exposition.

The suggested reading order is to start with Appendix A in the case of unfamiliarity with the calculus of mates, then read Section 1 and afterwards quickly become familiar with the statements in Appendix B, should this not already be the case. Then the reading of the main exposition in Sections 2 to 5 can begin, with some thumbing forwards to the statements in Appendix C and the statements and proofs in Appendix D when they are referenced. The proofs in Appendix C are there for completeness and only when the author was not able to find a good reference; the statements being quite basic, their proofs should not be considered a main part of this thesis and reading them is not necessary for understanding the central exposition.

Acknowledgments

First and foremost I would like to thank my advisor, Emanuele Dotto. He suggested the topic of this thesis, which I had a lot of fun working on, and guided me through the process of writing it, always being available when I had questions. I am also grateful for his many helpful suggestions for improving an earlier draft of this thesis.

Secondly I would like to thank Tashi Walde for a great number of very helpful discussions, which were invaluable in furthering my understanding of higher category theory, as well as proofreading parts of an earlier draft and helping to improve the introduction.

Furthermore I would like to thank Markus Land for providing me with a proof of a technical ∞ -categorical lemma that was needed for the construction of the Taylor graph.

Moreover I would like to thank Kevin Carlson, Denis Nardin, and Dylan Wilson for answering questions of mine on MathOverflow that occurred while working on this thesis. I am particularly grateful to Kevin Carlson for making me aware of the calculus of mates and the work of Riehl and Verity as well as the usefulness of their interplay.

Last but not least I would like to thank Emily Riehl for answering a question of mine via e-mail, going so far as to add the relevant statement to an ongoing book project.

1. Notation and conventions

When working with $(\infty, 1)$ -categories (which will be the case most of the time) we will use the framework of quasi-categories developed by Joyal, Lurie, and others. In particular, when we say ∞ -category we mean quasi-category. However, we will often work purely in the homotopy 2-category of ∞ -categories, thereby employing the theory developed by Riehl and Verity in a series of articles starting with [RV15] and their (as of September 2019) ongoing book project [RV]. Due to this, most of our arguments should not, in any fundamental way, depend on the precise model chosen for ∞ -categories.

Moreover, in quite a few places we will employ the calculus of mates of natural transformations. The needed facts are recalled in Appendix A, for those unfamiliar with the theory.

In the following we state the conventions and notations we will use throughout this thesis. A few basic facts concerning those notions that will be used throughout this thesis (often without explicit mention) are collected in Appendix B. Readers that are not well versed in the theory are advised to quickly familiarize themselves with the statements after finishing this first section.

Generally, if there is a pair of dual definitions or statements, we will only give one of them and leave the other implicit.

Convention 1.1. To avoid set-theoretic problems we will throughout assume that there is a sufficient supply of Grothendieck universes, so that any constructions we may employ will make sense in a potentially higher universe (this is the same approach as taken by Lurie, see [LurHTT, Section 1.2.15]). Objects belonging to the first such universe will be called *small*.

We will not assume ∞ -categories to be small, so that the examples we are interested in (such as the ∞ -category of all (small) spaces) are actually examples. Consequently we will not assume simplicial sets nor categories to be small (nor locally small), so that an ∞ -category is a simplicial set and its homotopy category a category. Posets, however, will be assumed to be small.

Convention 1.2. We will implicitly treat posets as categories and categories as ∞ -categories whenever it is convenient, without a change of notation. In particular, we will often just write “functor” for an order preserving map between posets.

Notation 1.3. We write **Pos** for the category of posets, **Pos**_∅ for the subcategory of posets that have an initial object together with initial object preserving functors, and **Pos**_∐ for the subcategory of posets that admit all (small) coproducts together with functors that preserve (small) coproducts.

Remark 1.4. Note that requiring a functor $f: \mathcal{P} \rightarrow \mathcal{Q}$ of posets to preserve I -indexed coproducts is equivalent to requiring equalities $f(\coprod_{i \in I} p_i) = \coprod_{i \in I} f(p_i)$. (For one direction we note that if the canonical map is an isomorphism, then it is already the identity; for the other that if we have the above equality, then the canonical map must be the identity since there are no other endomorphisms.)

Notation 1.5. We say a functor between categories is a *homotopy equivalence* if the geometric realization of its nerve is a (weak) homotopy equivalence. Similarly, we say a category is *contractible* if the geometric realization of its nerve is (weakly) contractible.

Notation 1.6. For \mathcal{I} a category we denote by $\mathcal{I}_{>\emptyset} \subseteq \mathcal{I}$ the full subcategory spanned by the non-initial objects.

Notation 1.7. We denote by $*$ the terminal category and, for an ∞ -category \mathcal{C} and $c \in \mathcal{C}$, by $\text{const}_c: * \rightarrow \mathcal{C}$ the functor representing c (sometimes we will also just write c for const_c). More generally, for a simplicial set K , we denote by $\text{const}_c: K \rightarrow \mathcal{C}$ the unique map that factors over $\text{const}_c: * \rightarrow \mathcal{C}$, and omit the index if the target category is $*$.

Notation 1.8. When \mathcal{I} , \mathcal{I}' , and \mathcal{J} are categories and $f: \mathcal{I} \rightarrow \mathcal{J}$ and $f': \mathcal{I}' \rightarrow \mathcal{J}$ are functors, we write $f \downarrow f'$ for the comma category and denote its objects by tuples $(i, i', f(i) \rightarrow f'(i'))$ where $i \in \mathcal{I}$ and $i' \in \mathcal{I}'$. Furthermore, we denote by $\text{pr}_{\mathcal{I}}$ respectively $\text{pr}_{\mathcal{I}'}$ (or just by pr (or $\text{pr}_{f \downarrow f'}$) if it is clear which one is meant) the forgetful functor from $f \downarrow f'$ to \mathcal{I} respectively \mathcal{I}' . Sometimes we will replace one or both of f and f' either with \mathcal{J} , in which case we mean the functor $\text{id}_{\mathcal{J}}$, or with an object $j \in \mathcal{J}$, in which case we mean the functor $\text{const}_j: * \rightarrow \mathcal{J}$. In the latter case, we will, when writing an object of the comma category, omit the unique object of $*$ from the tuple.

Remark 1.9. When \mathcal{J} is a poset, this comma category $f \downarrow f'$ is canonically isomorphic to the full subcategory of $\mathcal{I} \times \mathcal{I}'$ spanned by those (i, i') such that $f(i) \leq f'(i')$ (since any diagram in a poset automatically commutes). In particular we can omit mention of the structure maps $f(i) \rightarrow f'(i')$ in this case.

Notation 1.10. For \mathcal{C} an ∞ -category, we denote by $\text{h}\mathcal{C}$ its homotopy category (as a (1-)category, cf. [LurHTT, Section 1.2.3]) and by $\pi_c: \mathcal{C} \rightarrow \text{h}\mathcal{C}$ the canonical functor.

Notation 1.11. For K a simplicial set, we denote by K^\triangleleft the cone over K , by \blacktriangleleft the cone point, and by $\text{inc}_K: K \rightarrow K^\triangleleft$ the inclusion (we will sometimes drop the index if there is no risk of confusion).

Notation 1.12. For K a simplicial set and \mathcal{C} an ∞ -category, we will denote by $\text{Fun}(K, \mathcal{C})$ the ∞ -category of functors from K to \mathcal{C} , i.e. the internal hom of simplicial sets.

We will often implicitly identify $\text{Fun}(*, \mathcal{C})$ with \mathcal{C} itself.

Notation 1.13. Let $f: I \rightarrow J$ be a map of simplicial sets and \mathcal{C} an ∞ -category. We denote by $\text{Res}_f: \text{Fun}(J, \mathcal{C}) \rightarrow \text{Fun}(I, \mathcal{C})$ the restriction along f .

Notation 1.14. By an *adjunction* of functors between ∞ -categories we will mean an adjunction in the *homotopy 2-category of ∞ -categories*, i.e. the strict 2-category with objects the ∞ -categories, morphisms the functors between ∞ -categories, and 2-morphisms the homotopy classes of natural transformations between those functors (cf. [RV, Definition 1.4.1]).

Remark 1.15. This is the definition of an adjunction used by Riehl and Verity (see [RV, Definition 2.1.1]). We chose it since it is very pleasant to work with, in particular in relation to functors of Kan extension. That it agrees with the more hands-on definition of Lurie given in [LurHTT, Definition 5.2.2.1] is shown in [RV, Appendix F.5].

Definition 1.16. Let $f: I \rightarrow J$ be a map of simplicial sets. An ∞ -category \mathcal{C} is *weakly left f -extensible* if the restriction $\text{Res}_f: \text{Fun}(J, \mathcal{C}) \rightarrow \text{Fun}(I, \mathcal{C})$ has a left adjoint. In this case we fix such an adjunction $\text{Lan}_f \dashv \text{Res}_f$. In particular, we fix a unit-counit pair of this adjunction, which will be what we mean when we write “the” unit (or counit) of the adjunction.

Remark 1.17. If $f = \text{id}_I: I \rightarrow I$, then $\text{Res}_{\text{id}} = \text{id}$. In particular we can choose Lan_{id} , as well as the unit and counit of the adjunction, to be identities as well. This is the adjunction we fix in this case.

Notation 1.18. Let I be a simplicial set and \mathcal{C} an ∞ -category. Then we write $\Delta: \mathcal{C} \rightarrow \text{Fun}(I, \mathcal{C})$ for the diagonal, i.e. the restriction along $\text{const}: I \rightarrow *$, and say that \mathcal{C} *admits all colimits indexed by I* if it is weakly left const -extensible, i.e. if the functor Δ admits a left adjoint. In this case we write $\text{colim}_I := \text{Lan}_{\text{const}}: \text{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}$. Note that in particular we have a fixed adjunction $\text{colim}_I \dashv \text{Res}_{\text{const}}$.

Definition 1.19. Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories. We say that an ∞ -category \mathcal{C} is *left f -extensible* if it admits colimits indexed by $f \downarrow j$ for all $j \in \mathcal{J}$.

Remark 1.20. By [RV, Theorem 19.7.1, Lemma 13.5.1 (for $D = *$), Theorem 16.2.9 (iii), and Lemma 2.3.6], an ∞ -category \mathcal{C} that is left f -extensible is weakly left f -extensible (note that the proof of Theorem 19.7.1 does not need the whole completeness condition, only the weaker one we stated).

Moreover, the same sequence of statements (minus Lemma 2.3.6) implies that \mathcal{C} fulfills the property stated after [RV, Definition 13.5.7] which is what is needed in the proof of [RV, Proposition 13.5.8]. In particular we can apply this proposition whenever \mathcal{C} is left extensible with respect to the correct functors. This will be done a few times throughout Appendix C.

Remark 1.21. Note that an ∞ -category \mathcal{C} that admits colimits indexed by a category \mathcal{I} is left inc -extensible, where $\text{inc}: \mathcal{I} \rightarrow \mathcal{I}^\triangleright$ is the inclusion. This follows by considering, for an $a \in \mathcal{I}^\triangleright$, the slice category $\text{inc} \downarrow a$. If $a = \blacktriangleright$, it is isomorphic to \mathcal{I} . If otherwise $a \in \mathcal{I}$, it has a terminal object, in which case Lemma C.8 implies that \mathcal{C} admits colimits indexed by $\text{inc} \downarrow a$.

Notation 1.22. Let I be a simplicial set, \mathcal{C} an ∞ -category, and $p: I \rightarrow \mathcal{C}$ a diagram. We say that a diagram $I^\triangleright \rightarrow \mathcal{C}$ is a *colimit diagram extending p* if it is an initial object of $\mathcal{C}_{p/}$ (cf. [LurHTT, Remark 1.2.13.5]).

Notation 1.23. Let \mathcal{I} be a category, $i \in \mathcal{I}$, and \mathcal{C} an ∞ -category that admits colimits indexed by \mathcal{I} . Denote by $t_i: \Delta^1 \rightarrow \mathcal{I}^\triangleright$ the functor representing the unique morphism $i \rightarrow \blacktriangleright$. Then the functor $\text{Res}_{t_i}: \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C}) \rightarrow \text{Fun}(\Delta^1, \mathcal{C})$ carries to a natural transformation

$\alpha: \text{Res}_i \rightarrow \text{Res}_\blacktriangleright$ of functors $\text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) \rightarrow \mathcal{C}$. Now we can form the following composition (where the first equivalence is provided by Lemma B.2 and the last one by Lemma C.7)

$$\text{Res}_i \xrightarrow{\simeq} \text{Res}_i \text{Res}_{\text{inc}} \text{Lan}_{\text{inc}} = \text{Res}_i \text{Lan}_{\text{inc}} \xrightarrow{\alpha} \text{Res}_\blacktriangleright \text{Lan}_{\text{inc}} \xleftarrow{\simeq} \text{colim}_{\mathcal{I}}$$

which we consider to be the *structure map* from the value at i to the colimit.

Remark 1.24. We could have defined the structure map as the restriction of the unit $\text{id} \rightarrow \Delta \circ \text{colim}_{\mathcal{I}}$ to i , but the definition we gave is easier to compare to the notions of [LurHTT]. In particular note that, by Lemma B.3, the structure map $D(i) \rightarrow \text{colim}_{\mathcal{I}} D$ is an equivalence if and only if all colimit diagrams $\mathcal{T}^\triangleright \rightarrow \mathcal{C}$ extending D send the unique morphism $i \rightarrow \blacktriangleright$ to an equivalence (or, equivalently, if there exists one that does so).

Notation 1.25. Let \mathcal{I} be a category and \mathcal{C} be an ∞ -category that admits colimits indexed by \mathcal{I} . Then we have the following diagram on the left and its image under $\text{Fun}(-, \mathcal{C})$ on the right

$$\begin{array}{ccc} \mathcal{I} & \xrightarrow{\text{inc}} & \mathcal{T}^\triangleright \\ \downarrow & \xi \swarrow & \downarrow \text{id} \\ * & \blacktriangleright & \mathcal{T}^\triangleright \end{array} \qquad \begin{array}{ccc} \text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) & \xrightarrow{\text{Res}_{\text{id}}} & \text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) \\ \text{Res}_\blacktriangleright \downarrow & \xi \swarrow & \downarrow \text{Res}_{\text{inc}} \\ \mathcal{C} & \xrightarrow{\Delta} & \text{Fun}(\mathcal{I}, \mathcal{C}) \end{array}$$

where ξ is, at an object $i \in \mathcal{I}$, the unique map to the cone point. We obtain a mate $\xi_1: \text{colim}_{\mathcal{I}} \text{Res}_{\text{inc}} \rightarrow \text{Res}_\blacktriangleright \text{Lan}_{\text{id}} = \text{Res}_\blacktriangleright$. Evaluated at a diagram $D: \mathcal{T}^\triangleright \rightarrow \mathcal{C}$ this takes the form of a map

$$\text{colim}_{\mathcal{I}} D|_{\mathcal{I}} \longrightarrow D(\blacktriangleright)$$

that is natural in D . This is what we will mean by the *canonical map* from the colimit.

Remark 1.26. When $\mathcal{I} = *$, then $\text{colim}_{\mathcal{I}} = \text{id}$ and the natural transformation $\text{Res}_{\text{inc}} \rightarrow \text{Res}_\blacktriangleright$ is just given by ξ , i.e. evaluation at the unique morphism $* \rightarrow \blacktriangleright$ in $*^\triangleright$.

Notation 1.27. Let \mathcal{I} be a category with an initial object \emptyset , \mathcal{C} an ∞ -category that admits colimits indexed by $\mathcal{I}_{>\emptyset}$, and $D: \mathcal{I} \rightarrow \mathcal{C}$ a diagram. Then, noting that the full subcategory of \mathcal{I} spanned by \emptyset and $\mathcal{I}_{>\emptyset}$ is canonically isomorphic to $(\mathcal{I}_{>\emptyset})^\triangleleft$, we obtain, by the dual of what we did in Notation 1.25, a transformation $\text{Res}_\emptyset \rightarrow \lim_{\mathcal{I}_{>\emptyset}} \text{Res}_{\mathcal{I}_{>\emptyset}}$, i.e. a natural map

$$D(\emptyset) \longrightarrow \lim_{\mathcal{I}_{>\emptyset}} D|_{\mathcal{I}_{>\emptyset}}$$

which we see as the *canonical map* in this situation.

Notation 1.28. Let $f: I \rightarrow J$ be a map of simplicial sets and \mathcal{C} an ∞ -category that admits colimits indexed both by I and by J . Then we have the following diagram on the

left and its image under $\text{Fun}(-, \mathcal{C})$ on the right

$$\begin{array}{ccc}
I & \xrightarrow{f} & J \\
\downarrow & \text{id} \swarrow & \downarrow \\
* & \longrightarrow & *
\end{array}
\qquad
\begin{array}{ccc}
\mathcal{C} & \xrightarrow{\Delta_J} & \text{Fun}(J, \mathcal{C}) \\
\text{id} \downarrow & \text{id} \swarrow & \downarrow \text{Res}_f \\
\mathcal{C} & \xrightarrow{\Delta_I} & \text{Fun}(I, \mathcal{C})
\end{array}$$

whose mate $\text{id}_!$ is a natural transformation $\text{colim}_I \text{Res}_f \rightarrow \text{colim}_J$ which we will denote by f_* and call the *induced map* on the colimit.

Notation 1.29. We will say a map $f: K \rightarrow L$ of simplicial sets is *homotopy terminal* if, for each ∞ -category \mathcal{C} and colimit diagram $p: L^\triangleright \rightarrow \mathcal{C}$, the induced map $p \circ f^\triangleright: K^\triangleright \rightarrow \mathcal{C}$ is again a colimit diagram. The dual concept will be called *homotopy initial*.

Remark 1.30. By [LurHTT, Proposition 4.1.1.8], this definition of homotopy terminal is an equivalent characterization of what Lurie calls cofinal (cf. [LurHTT, Definition 4.1.1.1]).

Notation 1.31. Let $f: I \rightarrow J$ be a map of simplicial sets and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor between weakly left f -extensible ∞ -categories. We say that F *preserves left Kan extension along f* if the mate $\chi: \text{Lan}_f \circ (F \circ) \rightarrow (F \circ) \circ \text{Lan}_f$ of the natural transformation

$$\begin{array}{ccc}
\text{Fun}(J, \mathcal{C}) & \xrightarrow{\text{Res}_f} & \text{Fun}(I, \mathcal{C}) \\
F \circ \downarrow & \text{id} \swarrow & \downarrow F \circ \\
\text{Fun}(J, \mathcal{D}) & \xrightarrow{\text{Res}_f} & \text{Fun}(I, \mathcal{D})
\end{array}$$

is an equivalence.

We say that F *preserves colimits* indexed by a simplicial set I if it preserves left Kan extension along $\text{const}: I \rightarrow *$.

2. Excisive functors

Classical Goodwille calculus (as developed originally in the series of papers [Goo90; Goo92; Goo03]) and generalized to the ∞ -categorical context in [LurHA, Section 6]) studies functors which have certain behaviors with respect to diagrams indexed by cubes, i.e. posets of the following form:

Notation 2.1. Let $n \in \mathbb{N}_0$. We write $\mathcal{P}(n)$ for the n -cube, i.e. the poset of subsets of $[n-1] = \{0, \dots, n-1\}$ ordered by inclusion.

Namely, one defines a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between sufficiently nice ∞ -categories to be n -excisive if it sends strongly cocartesian $(n+1)$ -cubes to cartesian cubes. Here, cartesian means that the cube is a limit diagram, which makes sense in much greater generality:

Definition 2.2. Let \mathcal{I} be a category that has an initial object and \mathcal{C} an ∞ -category that admits limits indexed by $\mathcal{I}_{>\emptyset}$. A diagram $D: \mathcal{I} \rightarrow \mathcal{C}$ is *cartesian* if there is an initial object $\emptyset \in \mathcal{I}$ such the canonical map

$$D(\emptyset) \longrightarrow (\lim_{\mathcal{I}_{>\emptyset}} \text{Res}_{\mathcal{I}_{>\emptyset}})(D)$$

is an equivalence (by naturality of the map to the limit this is equivalent to requiring it to be an equivalence for each initial object).

Remark 2.3. By Lemma C.11, this is equivalent to requiring the restriction $D|_{\mathcal{J}}$ of D to the full subcategory \mathcal{J} of \mathcal{I} spanned by \emptyset and $\mathcal{I}_{>\emptyset}$ (which is canonically isomorphic to $(\mathcal{I}_{>\emptyset})^{\triangleleft}$) to fulfill the condition that the unit map $D|_{\mathcal{J}} \rightarrow (\text{Ran}_{\text{inc}} \text{Res}_{\text{inc}})(D|_{\mathcal{J}})$ is an equivalence, where $\text{inc}: \mathcal{I}_{>\emptyset} \rightarrow \mathcal{J}$ is the inclusion. By Lemmas C.1 and B.3, this is in turn equivalent to $D|_{\mathcal{J}}$ being a limit diagram. The latter description makes sense even if not all limits indexed by $\mathcal{I}_{>\emptyset}$ exist, which makes it useful in some circumstances.

Strongly cocartesian is a slightly more complicated condition: it means that any 2-face of the cube is cocartesian, i.e. a pushout (cf. [Goo92, Definition 2.1]). However, this can be rephrased in a more abstract way: it is equivalent to requiring the cube to be a left Kan extension of its restriction to the initial star, i.e. the following subposet:

Notation 2.4. Let $n \in \mathbb{N}_0$. We write $\mathcal{P}_{\leq 1}(n) \subseteq \mathcal{P}(n)$ for the full subposet consisting of all subsets of $[n-1]$ with cardinality at most 1 and denote the inclusion by \llcorner_n .

This rephrased condition does not make reference to the structure of the cube anymore, only to the inclusion \llcorner_n . In particular, we can formulate it for arbitrary functors:

Definition 2.5. Let $f: \check{\mathcal{I}} \rightarrow \mathcal{I}$ be a functor between categories and \mathcal{C} a weakly left f -extensible ∞ -category. A diagram $D: \mathcal{I} \rightarrow \mathcal{C}$ is *f -cocartesian* if the counit map $(\text{Lan}_f \text{Res}_f)(D) \rightarrow D$ is an equivalence.

Remark 2.6. That for $f = \llcorner_n$ this actually specializes to the condition of being a strongly cocartesian n -cube is shown in [LurHA, Proposition 6.1.1.15] and in slightly different language in [Goo92, Proposition 2.2].

Remark 2.7. If f is fully faithful, any diagram in the essential image of Lan_f will be f -cocartesian, by Lemma C.1.

Now that we have general notions of being “strongly cocartesian” with respect to some functor, we obtain a corresponding notion of excision for each of them:

Definition 2.8. Let $f: \check{\mathcal{I}} \rightarrow \mathcal{I}$ be a functor between categories such that \mathcal{I} has an initial object, \mathcal{C} a left f -extensible ∞ -category, and \mathcal{D} an ∞ -category admitting limits indexed by $\mathcal{I}_{>\emptyset}$. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is f -excisive if it takes f -cocartesian diagrams to cartesian diagrams. We write $\text{Exc}_f(\mathcal{C}, \mathcal{D}) \subseteq \text{Fun}(\mathcal{C}, \mathcal{D})$ for the full subcategory of f -excisive functors.

Moreover, for $n \in \mathbb{N}_0$, we say that a functor is n -excisive if it is \llcorner_{n+1} -excisive.

Remark 2.9. By Remark 2.6, our definition of an n -excisive functor agrees with Goodwillie’s original one in [Goo92, Definition 3.1]. In particular our notion of an f -excisive functor generalizes the classical one.

Our goal in this section is to show that under some hypotheses on a functor $f: \check{\mathcal{I}} \rightarrow \mathcal{I}$ and the ∞ -categories \mathcal{C} and \mathcal{D} there is, as in classical Goodwillie calculus, for any functor $F: \mathcal{C} \rightarrow \mathcal{D}$, a universal f -excisive functor approximating F which can be explicitly constructed.

For this, and the rest of this thesis, we will focus on functors of the following form, for reasons that will become apparent later.

Definition 2.10. A *preshape* is a functor $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ between posets such that \mathcal{S} has an initial object \emptyset and $\sigma^{-1}(\emptyset)$ is non-empty.

Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ and $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ be preshapes. A *map of preshapes* $\sigma \rightarrow \tau$ is a tuple (f, \check{f}) consisting of functors $f: \mathcal{S} \rightarrow \mathcal{T}$ and $\check{f}: \check{\mathcal{S}} \rightarrow \check{\mathcal{T}}$ such that $f \circ \sigma = \tau \circ \check{f}$ and $f^{-1}(\emptyset_{\mathcal{T}}) = \{\emptyset_{\mathcal{S}}\}$.

Remark 2.11. The collection of preshapes together with maps of preshapes forms a category with composition given by componentwise composition of functors.

Definition 2.12. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape.

- a) It is *finite* if both $\check{\mathcal{S}}$ and \mathcal{S} are finite.
- b) It is *full* if σ is a full functor.
- c) It is *reduced* if $\check{\mathcal{S}}$ has an initial object.

Remark 2.13. Since functors between posets are automatically faithful, a full preshape is already fully faithful. Furthermore, by Lemma D.1, it is also injective (on objects).

Remark 2.14. Note that a reduced preshape preserves the initial object (as $\sigma^{-1}(\emptyset_{\mathcal{S}})$ is downward closed). Furthermore, a full preshape is automatically reduced since fully faithful functors reflect initial objects.

Notation 2.15. When $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ is a reduced preshape, we denote the initial object of $\check{\mathcal{S}}$ by 0 .

Our construction of the universal excisive approximation is a generalization of Goodwillie's original construction for topological spaces [Goo03, Section 1] (which is also used in generalized form by Lurie in [LurHA, Section 6.1.1]). There, an important part is played by the cubical diagrams given by mapping a subset $U \subseteq [n-1]$ to the join of X and U , where X is some space and U is considered as a discrete space (one concrete construction of this join is to take, for each element of U , a copy of the cone of X and glue them all together at their bases). We will now describe a more abstract way of constructing these diagrams (which is basically the same way Lurie does it).

Notation 2.16. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced preshape and \mathcal{C} an ∞ -category with a terminal object. We write $\text{Pad}_{\check{\mathcal{S}}} := \text{Ran}_{\text{inc}}: \mathcal{C} \rightarrow \text{Fun}(\check{\mathcal{S}}, \mathcal{C})$, where $\text{inc}: \{0\} \rightarrow \check{\mathcal{S}}$ is the inclusion. Note that the categories $\check{s} \downarrow \text{inc}$ are, for all $\check{s} \in \check{\mathcal{S}}$, isomorphic to either the empty or the terminal category, and hence our assumption on \mathcal{C} imply the existence of this functor.

Remark 2.17. The diagram $\text{Pad}_{\check{\mathcal{S}}} X$ has X at 0 and the terminal object of \mathcal{C} at all other points of $\check{\mathcal{S}}$ (i.e. it pads the diagram with terminal objects).

Notation 2.18. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced preshape and \mathcal{C} a weakly left σ -extensible ∞ -category with a terminal object. Then the composition

$$\mathcal{C} \xrightarrow{\text{Pad}_{\check{\mathcal{S}}}} \text{Fun}(\check{\mathcal{S}}, \mathcal{C}) \xrightarrow{\text{Lan}_{\sigma}} \text{Fun}(\mathcal{S}, \mathcal{C})$$

curries to a functor $\mathcal{C} \times \mathcal{S} \rightarrow \mathcal{C}$ which we denote by \star .

Remark 2.19. By Lemma C.5, we have, for $X \in \mathcal{C}$ and $s \in \mathcal{S}$, the following more explicit formula

$$X \star s = (\text{Lan}_{\sigma} \text{Pad}_{\check{\mathcal{S}}})(X)(s) \simeq \text{colim}_{\sigma \downarrow s} (\text{Pad}_{\check{\mathcal{S}}}(X) \circ \text{pr}_{\sigma \downarrow s})$$

(as long as \mathcal{C} is left σ -extensible).

Remark 2.20. By definition of $\text{Pad}_{\check{\mathcal{S}}}(X)$, it admits a canonical map from any diagram $\check{\mathcal{S}} \rightarrow \mathcal{C}$ with X at 0. This is its main useful property and will allow us to factor maps into it in a useful way.

Remark 2.21. In the case where $\sigma = \lrcorner_n$ and \mathcal{C} is the ∞ -category of spaces, the functor \star specializes to the join with a discrete set (we can imagine the terminal objects occurring in $\text{Pad}_{\mathcal{P}_{\leq 1}(n)} X$ to be the cone over X which are then glued together by taking a left Kan extension). Hence \star generalizes the cubical diagrams mentioned above.

The following basic property will be needed later.

Lemma 2.22. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced preshape, \mathcal{I} a contractible category, and \mathcal{C} a left σ -extensible ∞ -category that admits colimits indexed by \mathcal{I} and has a terminal object. Then, for any $s \in \mathcal{S}$, the functor $(-\star s) = \text{Res}_s \text{Lan}_{\sigma} \text{Pad}_{\check{\mathcal{S}}}$ preserves terminal objects and colimits indexed by \mathcal{I} .*

Proof. Note that, by Lemma C.17, the functor Lan_σ preserves colimits and the functor $\text{Pad}_{\check{\mathcal{S}}}$ preserves limits. Hence, since Res_s also preserves colimits, it is enough to show that $\text{Pad}_{\check{\mathcal{S}}}$ preserves colimits indexed by \mathcal{I} and that $\text{Res}_s \text{Lan}_\sigma$ preserves terminal objects.

For the second statement note that an object of $\text{Fun}(\check{\mathcal{S}}, \mathcal{C})$ is terminal if and only if it is pointwise terminal. In particular, any such terminal object is equivalent to the restriction $\text{const}_* \circ c$, where $\text{const}_*: * \rightarrow \mathcal{C}$ represents a terminal object of \mathcal{C} and $c: \check{\mathcal{S}} \rightarrow *$ is the constant map. Now, by Lemma C.5, the object $(\text{Res}_s \text{Lan}_\sigma)(\text{const}_* \circ c)$ can be computed by $\text{colim}_{\sigma \downarrow s} (\text{const}_* \circ c \circ \text{pr}_{\sigma \downarrow s})$. But this is the terminal object of \mathcal{C} since $c \circ \text{pr}_{\sigma \downarrow s}$ is homotopy terminal as $\sigma \downarrow s$ has an initial object and is thus contractible.

For the first statement it is, by Lemma C.14, enough to show that, for any $\check{s} \in \check{\mathcal{S}}$, the functor $\text{Res}_{\check{s}} \text{Pad}_{\check{\mathcal{S}}}$ preserves colimits indexed by \mathcal{I} . Note that, again by Lemma C.5, the functor $\text{Res}_{\check{s}} \text{Pad}_{\check{\mathcal{S}}}$ is equivalent to $\lim_{\check{s} \downarrow \text{inc}} \circ \text{Res}_{\text{pr}_{\check{s} \downarrow \text{inc}}}$, where inc denotes the inclusion of $\{0\}$ into $\check{\mathcal{S}}$. But $\check{s} \downarrow \text{inc}$ is either the terminal category, in which case $\lim_{\check{s} \downarrow \text{inc}} = \text{id}$ clearly preserves colimits, or empty. In the latter case $\lim_{\check{s} \downarrow \text{inc}}$ is the functor $\text{const}_*: * \rightarrow \mathcal{C}$ for some terminal object $* \in \mathcal{C}$. This preserves colimits indexed by the contractible category \mathcal{I} since the constantly terminal diagram $\text{const}_*: \mathcal{I} \rightarrow \mathcal{C}$ is a colimit diagram extending $\text{const}_*: \mathcal{I} \rightarrow \mathcal{C}$ by [LurHTT, Proposition 4.3.1.12] (together with [LurHTT, Proposition 2.4.1.5]). \square

Now we are ready to give the construction of the universal excisive approximation. Note, however, that it will only have the desired properties after assuming more conditions on σ and the ∞ -categories \mathcal{C} and \mathcal{D} .

Construction 2.23. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced preshape, \mathcal{C} a left σ -extensible ∞ -category with a terminal object, and \mathcal{D} an ∞ -category admitting sequential colimits and limits indexed by $\mathcal{S}_{>\emptyset}$. We write

$$\text{T}_\sigma: \text{Fun}(\mathcal{C}, \mathcal{D}) \longrightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$$

for the functor given by

$$F \longmapsto \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} .$$

There is a natural transformation of functors $\mathcal{C} \rightarrow \mathcal{C}$

$$\tau_\sigma: \text{id} \xleftarrow{\simeq} \text{Res}_0 \text{Pad}_{\check{\mathcal{S}}} \longrightarrow \text{Res}_0 \text{Res}_\sigma \text{Lan}_\sigma \text{Pad}_{\check{\mathcal{S}}} = \text{Res}_\emptyset \text{Lan}_\sigma \text{Pad}_{\check{\mathcal{S}}}$$

coming from the counit of the adjunction $\text{Res}_0 \dashv \text{Pad}_{\check{\mathcal{S}}}$ and the unit of $\text{Res}_\sigma \dashv \text{Lan}_\sigma$, the first of which is an equivalence since the inclusion $\{0\} \subseteq \check{\mathcal{S}}$ is fully faithful (later we will need to assume that σ is full precisely because we need τ_σ and thus the mentioned unit to be equivalences). We obtain a natural transformation $t_\sigma: \text{id} \rightarrow \text{T}_\sigma$ of functors $\text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ defined at $F \in \text{Fun}(\mathcal{C}, \mathcal{D})$ by the composition

$$\begin{array}{ccc} F \xrightarrow{F \circ \tau_\sigma} F \circ \text{Res}_\emptyset \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} & & \text{T}_\sigma(F) \\ & \parallel & \parallel \\ & \text{Res}_\emptyset \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \longrightarrow \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} & \end{array}$$

where the second morphism is the canonical map to the limit. Now, by Lemma D.9, the sequence of morphisms

$$\mathrm{id} \xrightarrow{t_\sigma} T_\sigma \xrightarrow{t_\sigma \circ T_\sigma} T_\sigma T_\sigma \xrightarrow{t_\sigma \circ T_\sigma \circ T_\sigma} \dots \quad (2.1)$$

defines a sequential diagram. By our conditions on \mathcal{D} , its colimit exists. We will denote it by P_σ and by $p_\sigma: \mathrm{id} \rightarrow P_\sigma$ the structure map to the colimit.

Remark 2.24. The following is a more explicit formula for computing T_σ

$$T_\sigma(F)(X) \simeq \lim_{s \in \mathcal{S}_{>\emptyset}} F(X \star s)$$

which (in a less general form) was Goodwillie's original definition (cf. [Goo03, Section 1]). In this form t_σ is the composition of $F(X) \rightarrow F(X \star \emptyset)$ and the canonical map into the limit (the first of which is an equivalence if σ is full).

Remark 2.25. The construction of T_σ (and thus the one of P_σ) is *not* functorial in σ . The problem is that, although a map $(f, \check{f}): (\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}) \rightarrow (\tau: \check{\mathcal{T}} \rightarrow \mathcal{T})$ of reduced preshapes induces maps $X \star_\sigma s \rightarrow X \star_\tau f(s)$ and a map $\lim_{t \in \mathcal{T}_{>\emptyset}} (X \star_\tau t) \rightarrow \lim_{s \in \mathcal{S}_{>\emptyset}} (X \star_\tau f(s))$, they do not combine into a map between T_σ and T_τ . However, there are functorial properties when restricted to certain subcategories. This is discussed in Section 3.

For this construction to work well, we will need to assume further conditions on the target ∞ -category \mathcal{D} , namely that the occurring sequential colimits are compatible with certain limits.

Definition 2.26. Let \mathcal{I} be a category. An ∞ -category \mathcal{D} is *\mathcal{I} -differentiable* if it admits sequential colimits as well as limits indexed by $\mathcal{I}_{>\emptyset}$, and taking sequential colimits preserves limits indexed by $\mathcal{I}_{>\emptyset}$.

It is *differentiable* if it admits sequential colimits as well as all finite limits, and taking sequential colimits preserves finite limits.

Remark 2.27. The notion of a differentiable ∞ -category was introduced by Lurie, see [LurHA, Definition 6.1.1.6].

Remark 2.28. Note that differentiable implies \mathcal{I} -differentiable for any finite category \mathcal{I} (i.e. a category with finitely many objects and morphisms).

Remark 2.29. By Lemma C.18, the condition that sequential colimits preserve limits indexed by $\mathcal{S}_{>\emptyset}$ is equivalent to requiring the functor $\lim_{\mathcal{S}_{>\emptyset}}$ to preserve sequential colimits.

Example 2.30. The following are examples of differentiable ∞ -categories:

- any ∞ -topos (see [LurHA, Example 6.1.1.8]), in particular the ∞ -category of spaces (see [LurHTT, Proposition 6.3.4.1]).
- any stable ∞ -category (cf. [LurHA, Definition 1.1.1.9]) that admits countable coproducts (see [LurHA, Example 6.1.1.7]).

- the ∞ -category \mathcal{C}_* of pointed objects (cf. [LurHTT, Definition 7.2.2.1]) in a differentiable ∞ -category \mathcal{C} . In particular this tells us that the ∞ -category of pointed spaces is differentiable.

To see this, note that \mathcal{C}_* is defined as the full subcategory of $\text{Fun}(\Delta^1, \mathcal{C})$ spanned by the maps $f: \Delta^1 \rightarrow \mathcal{C}$ with $f(0)$ a terminal object of \mathcal{C} . But this subcategory is closed under the formation, in $\text{Fun}(\Delta^1, \mathcal{C})$, of finite limits and sequential colimits (the latter fact uses Lemma C.13). Since fully faithful functors reflect (co)limits by [RV, Proposition 4.3.12.], this implies that \mathcal{C}_* is differentiable.

We can already prove some elementary properties of T_σ and P_σ that we will need later.

Lemma 2.31. *Let $\sigma: \tilde{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced preshape, \mathcal{C} a left σ -extensible ∞ -category with a terminal object, and \mathcal{D} an \mathcal{S} -differentiable ∞ -category.*

- If σ is full, then, for any σ -excisive functor $F: \mathcal{C} \rightarrow \mathcal{D}$, both $t_\sigma(F)$ and $p_\sigma(F)$ are equivalences.*
- Both T_σ and P_σ preserve limits indexed by $\mathcal{S}_{>\emptyset}$.*
- Both T_σ and P_σ preserve sequential colimits.*
- Let \mathcal{C}' be another left σ -extensible ∞ -category with a terminal object and $F: \mathcal{C}' \rightarrow \mathcal{D}$ a functor. Then the functors $P_\sigma \circ (F \circ)$ and $(P_\sigma(F) \circ)$ from $\text{Fun}(\mathcal{C}, \mathcal{C}')$ to $\text{Fun}(\mathcal{C}, \mathcal{D})$ are equivalent when restricted to the full subcategory consisting of those functors $G: \mathcal{C} \rightarrow \mathcal{C}'$ that preserve terminal objects and left Kan extension along σ .*

Proof.

- The map $t_\sigma(F)$ is the composition of $F \circ \tau_\sigma$, which is an equivalence when σ is full, and the canonical map to the limit. The latter map is an equivalence by construction since, for any $X \in \mathcal{C}$, the diagram $(\text{Lan}_\sigma \text{Pad}_{\tilde{\mathcal{S}}})(X)$ is σ -cocartesian by Lemma C.1 and hence $F \circ (\text{Lan}_\sigma \text{Pad}_{\tilde{\mathcal{S}}})(X)$ cartesian. Thus $p_\sigma(F)$ is also an equivalence by Lemma C.13 as each map in diagram (2.1) is an equivalence (using that $(T_\sigma)^n(F)$ is, by induction, equivalent to F and hence σ -excisive).
- It follows directly from Lemmas C.15 and C.17 that T_σ preserves limits indexed by $\mathcal{S}_{>\emptyset}$. This also implies that the functor $\mathbb{N}_0 \rightarrow \text{Fun}(\text{Fun}(\mathcal{C}, \mathcal{D}), \text{Fun}(\mathcal{C}, \mathcal{D}))$ described by diagram (2.1) sends each $n \in \mathbb{N}_0$ to a functor that preserves limits indexed by $\mathcal{S}_{>\emptyset}$. Hence, by Lemma C.14, the associated functor $\text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathbb{N}_0, \text{Fun}(\mathcal{C}, \mathcal{D}))$ preserves limits indexed by $\mathcal{S}_{>\emptyset}$. Now, since \mathcal{D} is \mathcal{S} -differentiable, the functor $\text{colim}: \text{Fun}(\mathbb{N}_0, \text{Fun}(\mathcal{C}, \mathcal{D})) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ preserves limits indexed by $\mathcal{S}_{>\emptyset}$ (the category $\text{Fun}(\mathcal{C}, \mathcal{D})$ is again differentiable by Lemmas C.2 and C.15). As the composition of these two functors is precisely P_σ , this implies that P_σ preserves these limits as well.
- This follows similarly to b) by noting that $\lim: \text{Fun}(\mathcal{S}_{>\emptyset}, \mathcal{D}) \rightarrow \mathcal{D}$ preserves sequential colimits when \mathcal{D} is \mathcal{S} -differentiable.

- d) Using (the dual of) Lemma C.16, we see that any such G preserves right Kan extensions along the inclusion $\text{inc}: \{0\} \rightarrow \check{\mathcal{S}}$ (using that, for all $\check{s} \in \check{\mathcal{S}}$, the comma categories $\check{s} \downarrow \text{inc}$ are either empty or the terminal category). This implies the corresponding statement for T_σ , i.e. that there is an equivalence $\alpha: (T_\sigma(F) \circ) \rightarrow T_\sigma \circ (F \circ)$ (when restricted to the subcategory). Furthermore, by two applications of Lemma C.20, we obtain that the diagram

$$\begin{array}{ccc}
 (F \circ) & \xrightarrow{t_\sigma(F)} & (T_\sigma(F) \circ) \\
 \text{id} \downarrow & & \downarrow \alpha \\
 (F \circ) & \xrightarrow{t_\sigma} & T_\sigma \circ (F \circ)
 \end{array} \tag{2.2}$$

commutes up to homotopy.

Now we can inductively define equivalences $\alpha_n: ((T_\sigma)^n(F) \circ) \rightarrow (T_\sigma)^n \circ (F \circ)$ by setting $\alpha_0 = \text{id}$ and

$$\alpha_{n+1}: ((T_\sigma)^{n+1}(F) \circ) \xrightarrow{\alpha} T_\sigma \circ ((T_\sigma)^n(F) \circ) \xrightarrow{\alpha_n} T_\sigma \circ (T_\sigma)^n \circ (F \circ).$$

Now note that, since the diagram

$$\begin{array}{ccc}
 ((T_\sigma)^n(F) \circ) & \xrightarrow{t_\sigma} & (T_\sigma((T_\sigma)^n(F) \circ)) \\
 \text{id} \downarrow & & \downarrow \alpha \\
 ((T_\sigma)^n(F) \circ) & \xrightarrow{t_\sigma} & T_\sigma \circ ((T_\sigma)^n(F) \circ) \\
 \alpha_n \downarrow & & \downarrow \alpha_n \\
 (T_\sigma)^n \circ (F \circ) & \xrightarrow{t_\sigma} & T_\sigma \circ (T_\sigma)^n \circ (F \circ)
 \end{array}$$

commutes up to homotopy (using that the upper square is a special case of diagram (2.2)), the α_n assemble into an equivalence from the sequential diagram defining $(P_\sigma(F) \circ)$ to the sequential diagram defining $P_\sigma \circ (F \circ)$ (using Lemma D.9). \square

Remark 2.32. Note that in the first part of the previous lemma we did not use the full strength of F being σ -excisive, only that it sends diagrams in the essential image of $\text{Lan}_\sigma \text{Pad}_{\check{s}}$ to cartesian diagrams. Since we will show that (under certain conditions) the functor $P_\sigma F$ is always σ -excisive (see Lemma 2.46), this implies that (under these conditions) the a priori weaker property above is actually equivalent to being σ -excisive.

Sadly, it is not true that, for any preshape σ , the functor $P_\sigma(F)$ is a universal σ -excisive approximation to $F: \mathcal{C} \rightarrow \mathcal{D}$ (even if the ∞ -categories \mathcal{C} and \mathcal{D} are as nice as we want). This is shown by the following example:

Example 2.33. Consider the full subposet $\check{\mathcal{D}} := \{\emptyset, \{1, 2\}\} \subseteq \mathcal{P}(2)$ and denote the inclusion by δ . It is clear that δ is a full preshape. Now let \mathcal{C} and \mathcal{D} be two ∞ -categories. By

Lemma C.1, a diagram $\mathcal{P}(2) \rightarrow \mathcal{C}$ is δ -cocartesian if and only if it is equivalent to one of the form $\text{Lan}_\delta E$ for some diagram $E: \check{\mathcal{D}} \rightarrow \mathcal{C}$ (the diagram $\text{Lan}_\delta E$ looks like

$$\begin{array}{ccc} X & \xrightarrow{\text{id}} & X \\ \text{id} \downarrow & & \downarrow f \\ X & \xrightarrow{f} & Y \end{array}$$

where f is the morphism of \mathcal{C} represented by $E: \Delta^1 \cong \check{\mathcal{D}} \rightarrow \mathcal{C}$). In particular, for a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ to be δ -excisive it has to send any such $\text{Lan}_\delta E$ to a cartesian diagram. For Y the terminal object of \mathcal{C} this would imply, if F preserves terminal objects, that

$$F(X) \xleftarrow{\text{id}} F(X) \xrightarrow{\text{id}} F(X)$$

exhibits $F(X)$ as a product of $F(X)$ with itself. In the case where \mathcal{D} is the ∞ -category \mathbf{S}_* of pointed spaces, this can only be the case if $F(X)$ is weakly contractible (by considering its homotopy groups). This shows that a δ -excisive functor $\mathcal{C} \rightarrow \mathbf{S}_*$ that preserves terminal objects has its image contained in the terminal objects.

Now let $F: \mathcal{C} \rightarrow \mathbf{S}_*$ be any functor that preserves terminal objects. Since $\text{Pad}_{\check{\mathcal{D}}}$ is given by sending $X \in \mathcal{C}$ to a map $X \rightarrow *$, we obtain that $\text{T}_\delta(F)$ is the functor $X \mapsto F(X) \times F(X)$ and that t_δ is given, at X , by the diagonal $F(X) \rightarrow F(X) \times F(X)$. In particular, we have that $\text{P}_\delta(F)(X)$ is given by the colimit of the sequence

$$F(X) \xrightarrow{\Delta} F(X) \times F(X) \xrightarrow{\Delta} (F(X) \times F(X)) \times (F(X) \times F(X)) \longrightarrow \dots$$

and hence is not weakly contractible when $F(X)$ is not weakly contractible since homotopy groups commute with products and sequential (homotopy) colimits (for 1-categorical sequential colimits over inclusions this can be found in [May, Chapter 9.4]; for sequential homotopy colimits it follows from the 1-categorical case by taking a cofibrant replacement). Thus, if the image of F is not contained in the terminal objects, then $\text{P}_\delta(F)$ can not be δ -excisive.

However, we can put conditions on σ so that P_σ is a functor of universal σ -excisive approximation. This is done in the next definition. We will see that L_n fulfills these conditions, so that our statements actually generalize the classical ones.

Definition 2.34. A preshape $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ is a *shape* if \mathcal{S} has all (small) coproducts, and, for all $s, t \in \mathcal{S}$ and $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \leq t \amalg s$, the full subposet

$$\check{\mathcal{S}}_{s,t,\check{k}} := \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s \text{ and } \sigma(\check{k}) \leq t \amalg \sigma(\check{s})\} \subseteq \check{\mathcal{S}}$$

is contractible.

A *map of shapes* is a map between the underlying preshapes.

Remark 2.35. For everything we do with shapes in this section it would be enough to only require \mathcal{S} to admit finite coproducts. However, we use the stronger version since it makes the constructions in Section 4.2 easier to work with (though one should be able to work around this, so that in the end (almost) all statements we will make should also hold with the weaker requirements).

The following is an easy-to-check sufficient criterion for a reduced preshape to be a shape.

Lemma 2.36. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full preshape. Assume that, for all $a, b \in \mathcal{S}$ and $\check{c} \in \check{\mathcal{S}}$ such that $\sigma(\check{c}) \leq a \amalg b$, we have $\sigma(\check{c}) \leq a$ or $\sigma(\check{c}) \leq b$. Then σ is a shape.*

Proof. We want to show that, for all $s, t \in \mathcal{S}$ and $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \leq t \amalg s$, the poset $\check{\mathcal{S}}_{s,t,\check{k}}$ is contractible. For this first assume that $\sigma(\check{k}) \leq t$. In this case $\check{\mathcal{S}}_{s,t,\check{k}}$ has 0 as an initial object and is thus contractible. Otherwise $\sigma(\check{k}) \not\leq t$ and our assumption implies $\sigma(\check{k}) \leq s$ and hence $\check{k} \in \check{\mathcal{S}}_{s,t,\check{k}}$. Thus we have that $\sigma(\check{k}) \leq \sigma(\check{s})$ for any $\check{s} \in \check{\mathcal{S}}_{s,t,\check{k}}$. As σ is full this implies that \check{k} is an initial object of $\check{\mathcal{S}}_{s,t,\check{k}}$, which finishes the proof. \square

Example 2.37. We give some examples and non-examples for Definition 2.34:

- The preshape \perp_n is a shape. This follows directly from Lemma 2.36. (A more general version of this statement will be proven in Lemma 4.14.)
- More generally, let $\check{\mathcal{S}}$ be any non-empty downward closed full subposet of $\mathcal{P}(n)$. Then the inclusion $\check{\mathcal{S}} \rightarrow \mathcal{P}(n)$ is a shape. To see this, let $S, T \in \mathcal{P}(n)$ and $\check{K} \in \check{\mathcal{S}}$ such that $\check{K} \subseteq S \cup T$, and note that $\check{\mathcal{S}}_{S,T,\check{K}} = \{\check{S} \in \check{\mathcal{S}} \mid \check{K} \setminus T \subseteq \check{S} \subseteq S\}$. Now, since $\check{\mathcal{S}}$ is downward closed, the element $\check{K} \setminus T \in \mathcal{P}(n)$ is contained in $\check{\mathcal{S}}$, and hence is an initial object of $\check{\mathcal{S}}_{S,T,\check{K}}$.
- The preshape δ from Example 2.33 is not a shape. For example the poset $\check{\mathcal{D}}_{s,t,\check{k}}$ for $s = \{1\}$, $t = \{2\}$, and $\check{k} = \{1, 2\}$ is empty and thus not contractible.

The main motivation for the definition of a shape is that it is precisely what we need for the next lemma. However, before we can state it, we need a way to assume, depending on the shape, enough colimits to exist. This will be achieved by the following definition. It is somewhat stronger than what we will actually need, but a lot more convenient to work with (it would be possible to track the precise requirements; however we chose to not do so in favor of increased readability).

Definition 2.38. Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between posets and \mathcal{C} an ∞ -category. If both \mathcal{I} and \mathcal{J} are finite, we say that \mathcal{C} is *f-nice* if it admits all finite colimits. Otherwise we say that \mathcal{C} is *f-nice* if it admits colimits of size up to the maximum of the cardinalities of \mathcal{I} and \mathcal{J} .

Remark 2.39. Note that an *f-nice* ∞ -category is automatically left *g*-extensible for any functor *g* with source \mathcal{I} or \mathcal{J} and target a poset (since the corresponding slice categories have a cardinality bounded by the cardinality of \mathcal{I} respectively \mathcal{J}).

Lemma 2.40. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a shape, \mathcal{C} a σ -nice ∞ -category, and $D: \check{\mathcal{S}} \rightarrow \mathcal{C}$ a diagram. Then, for any $t \in \mathcal{S}$, the diagram $(\text{Lan}_\sigma D) \circ (t \amalg -)$ is σ -cocartesian. In particular $\text{Lan}_\sigma D$ is σ -cocartesian, even though σ is not necessarily full (in which case Lemma C.1 would imply the statement).*

This also implies that, for any σ -cocartesian diagram $D': \mathcal{S} \rightarrow \mathcal{C}$ and $t \in \mathcal{S}$, the diagram $D' \circ (t \amalg -)$ is again σ -cocartesian.

Proof. This is a special case of the (technical) next lemma. More precisely, we apply it to the situation

$$\begin{array}{ccc} & \check{\mathcal{S}} & \xrightarrow{D} \mathcal{C} \\ & \downarrow \sigma & \\ \check{\mathcal{S}} & \xrightarrow{\sigma} \mathcal{S} & \xrightarrow{t\Pi-} \mathcal{S} \end{array}$$

for which we need that, for all $s \in \mathcal{S}$ and $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \leq (t\Pi-)(s) = t\Pi s$, the poset

$$\{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s \text{ and } \sigma(\check{k}) \leq t\Pi\sigma(\check{s})\} = \check{\mathcal{S}}_{s,t,\check{k}}$$

is contractible. But this is precisely the assumption on σ for it to be a shape. \square

Lemma 2.41. *Let \mathcal{I} , \mathcal{J} , \mathcal{K} , and \mathcal{L} be posets, and f , g , and h functors as the following diagram*

$$\begin{array}{ccc} & \mathcal{I} & \\ & \downarrow f & \\ \mathcal{J} & \xrightarrow{g} \mathcal{K} \xrightarrow{h} & \mathcal{L} \end{array}$$

specifies and \mathcal{C} an f -nice and g -nice ∞ -category. Furthermore, assume that, for all $k \in \mathcal{K}$ and $i \in \mathcal{I}$ such that $f(i) \leq h(k)$, the full subposet

$$\{j \in \mathcal{J} \mid f(i) \leq h(g(j)) \text{ and } g(j) \leq k\} \subseteq \mathcal{J}$$

is contractible. Then, for any diagram $D: \mathcal{I} \rightarrow \mathcal{C}$, the diagram $(\text{Res}_h \text{Lan}_f)(D)$ is g -cocartesian.

Proof. Consider the following diagram

$$\begin{array}{ccc} f \downarrow (h \circ g) & \xrightarrow{p} & \mathcal{I} \\ \downarrow q & \swarrow \alpha & \downarrow f \\ \mathcal{J} & \xrightarrow{h \circ g} & \mathcal{L} \\ \downarrow g & \swarrow \text{id} & \downarrow \text{id} \\ \mathcal{K} & \xrightarrow{h} & \mathcal{L} \end{array}$$

where $p = \text{pr}_{\mathcal{I}}$ and $q = \text{pr}_{\mathcal{J}}$ are the two projections and α comes from $f(p(i, j)) = f(i) \leq h(g(j)) = h(g(q(i, j)))$. Applying $\text{Fun}(-, \mathcal{C})$ yields the following diagram on the left and

subsequently taking mates the one on the right

$$\begin{array}{ccc}
\text{Fun}(f \downarrow (h \circ g), \mathcal{C}) & \xleftarrow{\text{Res}_p} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
\uparrow \text{Res}_q & \swarrow \alpha & \uparrow \text{Res}_f \\
\text{Fun}(\mathcal{J}, \mathcal{C}) & \xleftarrow{\text{Res}_{h \circ g}} & \text{Fun}(\mathcal{L}, \mathcal{C}) \\
\uparrow \text{Res}_g & \swarrow \text{id} & \uparrow \text{Res}_{\text{id}} \\
\text{Fun}(\mathcal{K}, \mathcal{C}) & \xleftarrow{\text{Res}_h} & \text{Fun}(\mathcal{L}, \mathcal{C})
\end{array}
\qquad
\begin{array}{ccc}
\text{Fun}(f \downarrow (h \circ g), \mathcal{C}) & \xleftarrow{\text{Res}_p} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
\downarrow \text{Lan}_q & \swarrow \alpha_! & \downarrow \text{Lan}_f \\
\text{Fun}(\mathcal{J}, \mathcal{C}) & \xleftarrow{\text{Res}_{h \circ g}} & \text{Fun}(\mathcal{L}, \mathcal{C}) \\
\downarrow \text{Lan}_g & \swarrow \text{id}_! & \downarrow \text{Lan}_{\text{id}} \\
\text{Fun}(\mathcal{K}, \mathcal{C}) & \xleftarrow{\text{Res}_h} & \text{Fun}(\mathcal{L}, \mathcal{C})
\end{array}$$

(where all occurring Kan extensions exist by our assumptions on \mathcal{C}).

Now note that since, by definition, the mate $\text{id}_!$ is given by the composition

$$\text{Lan}_g \text{Res}_{h \circ g} = \text{Lan}_g \text{Res}_{h \circ g} \text{Res}_{\text{id}} \text{Lan}_{\text{id}} = \text{Lan}_g \text{Res}_g \text{Res}_h \text{Lan}_{\text{id}} \xrightarrow{\varepsilon} \text{Res}_h \text{Lan}_{\text{id}} = \text{Res}_h$$

applying $\text{id}_!$ to a diagram $D: \mathcal{L} \rightarrow \mathcal{C}$ results precisely in the map from the definition of g -cocartesianess for $\text{Res}_h(D)$. In particular it is enough to show that $\text{id}_! \circ \text{Lan}_f$ is an equivalence, which is one of the maps that occur in the paste $\text{id}_! * \alpha_! = (\text{id} * \alpha)_!$ (where the equality (in the homotopy 2-category of ∞ -categories) comes from the pasting law for mates). Since $\alpha_!$ is an equivalence by Lemma C.4, it is thus enough to show that $(\text{id} * \alpha)_!: \text{Lan}_{g \circ q} \text{Res}_p \rightarrow \text{Res}_h \text{Lan}_{\text{id} \circ f}$ is one as well.

For this it is, by Lemma C.6, enough to show that for any $k \in \mathcal{K}$, the map

$$\text{colim}_{(g \circ q) \downarrow k} \text{Res}_{\text{pr}_{(g \circ q) \downarrow k}} \text{Res}_p \xrightarrow{r_*} \text{colim}_{f \downarrow h(k)} \text{Res}_{\text{pr}_{f \downarrow h(k)}}$$

induced by the functor $r: (g \circ q) \downarrow k \rightarrow f \downarrow h(k)$ given by p is an equivalence. We claim that r is homotopy terminal, for which we need that, for all $i \in f \downarrow h(k)$, the poset

$$i \downarrow r = \{(i', j') \in \mathcal{I} \times \mathcal{J} \mid f(i') \leq h(g(j')) \text{ and } g(j') \leq k \text{ and } i \leq i'\}$$

is contractible. For this note that the map

$$\mathcal{Q} := \{j' \in \mathcal{J} \mid f(i) \leq h(g(j')) \text{ and } g(j') \leq k\} \rightarrow i \downarrow r$$

given by $j' \mapsto (i, j')$ is left adjoint to the projection $i \downarrow r \rightarrow \mathcal{Q}$ given by $(i', j') \mapsto j'$ and hence a homotopy equivalence. Thus, it is enough to show that the poset on the left is contractible, which is true by assumption. \square

We can now formulate the main result of this section.

Theorem 2.42. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, and \mathcal{D} an \mathcal{S} -differentiable ∞ -category. Then there is an adjunction with left adjoint $\text{P}_\sigma: \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Exc}_\sigma(\mathcal{C}, \mathcal{D})$, right adjoint the inclusion $\text{inc}: \text{Exc}_\sigma(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$, and unit $\text{p}_\sigma: \text{id} \rightarrow \text{inc} \circ \text{P}_\sigma$.*

Remark 2.43. Later we will also obtain Corollary 4.8, a version of this theorem for shapes which are finite but not necessarily full.

The main input in proving this theorem is the following lemma, whose proof is adapted from Rezk's streamlined proof (see [Rez]) of the corresponding statement for ordinary Goodwillie calculus.

Lemma 2.44. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, \mathcal{D} an ∞ -category admitting sequential colimits and limits indexed by $\mathcal{S}_{>\emptyset}$, $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor, and $D: \mathcal{S} \rightarrow \mathcal{C}$ a σ -cocartesian diagram. Then there is a homotopy commutative diagram*

$$\begin{array}{ccc} F \circ D & \xrightarrow{t_\sigma(F)} & T_\sigma(F) \circ D \\ & \searrow & \nearrow \\ & E & \end{array}$$

such that $E: \mathcal{S} \rightarrow \mathcal{D}$ is cartesian.

Proof. The general strategy is to define, dependent on D , the following data

- a functor $\mathcal{S} \times \mathcal{S} \rightarrow \mathcal{C}$ written $(s, t) \mapsto D_s(t)$ and $D': \mathcal{S} \rightarrow \text{Fun}(\mathcal{S}, \mathcal{C})$, $t \mapsto D_-(t)$ after currying,
- a natural transformation $\alpha: D' \rightarrow \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \circ D$ (which has the pointwise form $\alpha: D_s(t) \rightarrow D(t) \star s$),
- and a natural transformation $\beta: D \rightarrow \text{Res}_\emptyset \circ D' = D_\emptyset(-)$

such that

1. the composition $(\text{Res}_\emptyset \circ \alpha)\beta: D \rightarrow \text{Res}_\emptyset \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \circ D$ is homotopic to $\tau_\sigma \circ D$ (where τ_σ is as in Construction 2.23).
2. for each $s \in \mathcal{S}_{>\emptyset}$ the diagram $F \circ \text{Res}_s \circ D' = \text{Res}_s \circ (F \circ) \circ D': \mathcal{S} \rightarrow \mathcal{D}$, which is more explicitly given by $t \mapsto F(D_s(t))$, is cartesian.

Assuming this exists, we obtain the homotopy commutative diagram

$$\begin{array}{ccc} & & F \circ D \\ & \xleftarrow{\beta} & \downarrow \tau_\sigma \\ F \circ \text{Res}_\emptyset \circ D' & \xrightarrow{\alpha} & F \circ \text{Res}_\emptyset \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \circ D \\ \parallel & & \parallel \\ \text{Res}_\emptyset \circ (F \circ) \circ D' & \xrightarrow{\alpha} & \text{Res}_\emptyset \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \circ D \\ \downarrow & & \downarrow \\ \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ D' & \xrightarrow{\alpha} & \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} \circ D = T_\sigma(F) \circ D \end{array}$$

which proves the claim by setting $E := \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ D'$ and noting that E is cartesian by Lemma D.12 and the assumption that $\text{Res}_s \circ (F \circ) \circ D'$ is cartesian for each $s \in \mathcal{S}_{>\emptyset}$.

Now for the construction. Since it needs to work for arbitrary F , the easiest way to obtain cartesian diagrams $F \circ D_s(-)$ is to have the $D_s(-)$ be diagrams that are cartesian because they contain a lot of equivalences, which are preserved by any functor. Namely, let $s \in \mathcal{S}_{>\emptyset}$, and assume that, for all $t \in \mathcal{S}$, applying D_s to the morphism $t \rightarrow t \amalg s$ yields an equivalence. Then $F \circ D_s$ is cartesian by Lemma 2.45, which can be found after this proof (this is where we essentially need that \mathcal{S} is a poset).

This suggests to try to define $D_s(t) = D(t \amalg s)$, as then $(- \amalg s)$ being idempotent in a poset implies the needed condition. However, it is not clear how one should then define the natural transformation $D(t \amalg s) \rightarrow D(t) \star s$. So, our strategy is to construct $D_s(t)$ in a way that comes with this map but, in good cases, still computes $D(t \amalg s)$.

The rough idea is to set

$$D_s(t) := (\text{Lan}_\sigma \text{Res}_\sigma)(D \circ (t \amalg -))(s)$$

which will come equipped with a map $D_-(t) \rightarrow D(t) \star -$ since the latter is defined as $(\text{Lan}_\sigma \text{Pad}_{\check{\mathcal{S}}})(D(t))$ which is terminal in the correct sense (see Remark 2.20). Furthermore, for $D_-(t)$ to compute $D(t \amalg -)$, we precisely need that $D \circ (t \amalg -)$ is σ -cocartesian, which was the statement of Lemma 2.40.

More formally, to also obtain functoriality in t , we consider the diagram

$$\begin{array}{ccccccc} \mathcal{S} & \xrightarrow{p'} & \text{Fun}(\mathcal{S}, \mathcal{C}) & \xrightarrow{\text{Res}_\sigma} & \text{Fun}(\check{\mathcal{S}}, \mathcal{C}) & \xrightarrow{\text{id}} & \text{Fun}(\check{\mathcal{S}}, \mathcal{C}) & \xrightarrow{\text{Lan}_\sigma} & \text{Fun}(\mathcal{S}, \mathcal{C}) \\ & & & & \searrow & \Downarrow \eta & \nearrow & & \\ & & & & \text{Res}_0 & & \text{Pad}_{\check{\mathcal{S}}} & & \\ & & \searrow D & & & & & & \\ & & & & & & & & \mathcal{C} \end{array}$$

where p' is given by $t \mapsto D \circ (t \amalg -)$ and η is the unit of the adjunction $\text{Res}_0 \dashv \text{Pad}_{\check{\mathcal{S}}}$. The upper composition gives the desired functor $D': \mathcal{S} \rightarrow \text{Fun}(\mathcal{S}, \mathcal{C})$ and η the natural transformation α , using that the left triangle commutes as $t \amalg \sigma(0) = t$.

Furthermore, let β be given by the natural transformation in the diagram

$$\begin{array}{ccccc} \mathcal{S} & & & & \\ \downarrow p' & & & & \\ \text{Fun}(\mathcal{S}, \mathcal{C}) & & & & \\ \downarrow \text{Res}_\sigma & & & & \\ \text{Fun}(\check{\mathcal{S}}, \mathcal{C}) & \xrightarrow{\text{id}} & \text{Fun}(\check{\mathcal{S}}, \mathcal{C}) & \xrightarrow{\text{Res}_0} & \mathcal{C} \\ \downarrow \text{Lan}_\sigma & & \downarrow \varepsilon & \nearrow \text{Res}_\sigma & \\ \text{Fun}(\mathcal{S}, \mathcal{C}) & & & & \\ & \searrow \text{Res}_\sigma & & & \\ & & \text{Fun}(\mathcal{S}, \mathcal{C}) & \xrightarrow{\text{Res}_\sigma} & \mathcal{C} \end{array}$$

where ε is the unit of the adjunction $\text{Lan}_\sigma \dashv \text{Res}_\sigma$.

To show that $(\text{Res}_\emptyset \circ \alpha)\beta$ is homotopic to $\tau_\sigma \circ D$, write $p = \text{Res}_\sigma \circ p'$ and consider the diagram

$$\begin{array}{ccccc}
\text{Res}_\emptyset \text{Res}_\sigma \text{Lan}_\sigma p & \xleftarrow{\beta} & \text{Res}_\emptyset p & \xrightarrow{\text{id}} & \text{Res}_\emptyset p \\
\parallel & & \downarrow & & \uparrow \simeq \\
\text{Res}_\emptyset \text{Lan}_\sigma p & & \text{Res}_\emptyset(\text{Pad}_{\mathcal{S}} \text{Res}_\emptyset)p = (\text{Res}_\emptyset \text{Pad}_{\mathcal{S}}) \text{Res}_\emptyset p & & \\
\alpha \downarrow & & \downarrow & & \\
\text{Res}_\emptyset \text{Lan}_\sigma \text{Pad}_{\mathcal{S}} \text{Res}_\emptyset p = \text{Res}_\emptyset \text{Res}_\sigma \text{Lan}_\sigma \text{Pad}_{\mathcal{S}} \text{Res}_\emptyset p & & & &
\end{array}$$

where all maps (apart from the four identities) are given by the (co)units of the adjunctions $\text{Res}_\emptyset \dashv \text{Pad}_{\mathcal{S}}$ and $\text{Lan}_\sigma \dashv \text{Res}_\sigma$, and the right vertical morphism is an equivalence since the inclusion $\{0\} \subseteq \mathcal{S}$ is fully faithful. The right square commutes up to homotopy by one of the triangle identities and the left square commutes up to homotopy since the two sides are just the two possible horizontal compositions of a pair of natural transformations. Now note that the composition along the left of the diagram is just $(\text{Res}_\emptyset \circ \alpha)\beta$ and the one along the right is $\tau_\sigma \circ D$.

The only thing left to show is that $D_s(-) = \text{Res}_s D'$ is equivalent to $D(- \amalg s) = \text{Res}_s p'$. This then implies that the morphism $D_s(t) \rightarrow D_s(t \amalg s)$ is an equivalence for all $t \in \mathcal{S}$, as $t \amalg s \rightarrow (t \amalg s) \amalg s$ is the identity. We will actually even show that the map

$$D' = \text{Lan}_\sigma \text{Res}_\sigma p' \xrightarrow{\varepsilon} p'$$

is an equivalence, where ε is the counit of the adjunction $\text{Lan}_\sigma \dashv \text{Res}_\sigma$. For this it is enough to show that ε evaluated at $p'(t) = D \circ (t \amalg -)$ is an equivalence for all $t \in \mathcal{S}$. But this is equivalent to $D \circ (t \amalg -)$ being σ -cocartesian, which was the statement of Lemma 2.40. \square

Lemma 2.45. *Let \mathcal{I} be a poset that admits finite coproducts, \emptyset its initial object, i an element of $\mathcal{I}_{>\emptyset}$, \mathcal{C} an ∞ -category that admits limits indexed by $\mathcal{I}_{>\emptyset}$, and $D: \mathcal{I} \rightarrow \mathcal{C}$ a diagram. Furthermore, assume that D applied to the morphism $j \rightarrow j \amalg i$ is an equivalence for all $j \in \mathcal{I}$. Then D is cartesian.*

Proof. Let $p_i: \mathcal{I} \rightarrow i \downarrow \mathcal{I}_{>\emptyset}$ be the functor given by $j \mapsto j \amalg i$. Note that the restriction $p_i|_{\mathcal{I}_{>\emptyset}}$ is left adjoint to the projection $\text{pr}: i \downarrow \mathcal{I}_{>\emptyset} \rightarrow \mathcal{I}_{>\emptyset}$ by the universal property of the coproduct. Additionally, let $\eta: \text{id} \rightarrow \text{inc} \circ \text{pr} \circ p_i$ be the natural transformation of functors $\mathcal{I} \rightarrow \mathcal{I}$ coming from $j \leq j \amalg i$ (this restricts to the unit of the above adjunction on $\mathcal{I}_{>\emptyset}$).

Now consider the diagram $D': (i \downarrow \mathcal{I}_{>\emptyset})^\triangleleft \rightarrow \mathcal{C}$ given by the composition of $D \circ \text{inc} \circ \text{pr}$ and the functor $(i \downarrow \mathcal{I}_{>\emptyset})^\triangleleft \rightarrow i \downarrow \mathcal{I}_{>\emptyset}$ given by i at the cone point and the identity otherwise. From this, we obtain the following diagram

$$\begin{array}{ccccc}
D(\emptyset) & \xrightarrow{\simeq_\eta} & (D \circ \text{inc} \circ \text{pr} \circ p_i)(\emptyset) & \xlongequal{\quad} & (D \circ \text{inc} \circ \text{pr})(i) \\
\downarrow & & \downarrow & & \downarrow \simeq \\
\lim_{\mathcal{I}_{>\emptyset}} (D \circ \text{inc}) & \xrightarrow{\simeq_\eta} & \lim_{\mathcal{I}_{>\emptyset}} (D \circ \text{inc} \circ \text{pr} \circ p_i \circ \text{inc}) & \xleftarrow{\simeq_{(p_i \circ \text{inc})^*}} & \lim_{i \downarrow \mathcal{I}_{>\emptyset}} (D \circ \text{inc} \circ \text{pr})
\end{array}$$

which commutes up to homotopy by (the dual of) Lemma C.9 (note that, by Lemma C.8, limits indexed by the poset $i \downarrow \mathcal{I}_{>\emptyset}$ exist since it has an initial object). The left horizontal morphisms are equivalences as $D \circ \eta$ is an equivalence by assumption. Furthermore, the bottom right horizontal map is an equivalence since $p_i \circ \text{inc}$ is left adjoint and hence homotopy initial, and the vertical morphism on the right is an equivalence by Lemma C.10 since i is an initial object of $i \downarrow \mathcal{I}_{>\emptyset}$. Hence the vertical map on the left is an equivalence, which was the claim. \square

The rest of the proof of Theorem 2.42 is analogous to the one given by Lurie in [LurHA, 6.1.1.10]. However, for completeness' sake, we still recount it here (with a bit more details). We need a few more lemmas:

Lemma 2.46. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, and \mathcal{D} an \mathcal{S} -differentiable ∞ -category. Then, for any functor $F: \mathcal{C} \rightarrow \mathcal{D}$, the functor $P_\sigma(F): \mathcal{C} \rightarrow \mathcal{D}$ is σ -excisive.*

Proof. Let $D: \mathcal{S} \rightarrow \mathcal{C}$ be a σ -cocartesian diagram. By definition, the diagram $P_\sigma(F) \circ D$ is given by the colimit of

$$F \circ D \longrightarrow T_\sigma(F) \circ D \longrightarrow (T_\sigma)^2(F) \circ D \longrightarrow \dots \quad (2.3)$$

where the maps are given by t_σ . By Lemma 2.44, each of these maps

$$t_\sigma((T_\sigma)^n(F)) \circ D: (T_\sigma)^n(F) \circ D \longrightarrow (T_\sigma)^{n+1}(F) \circ D$$

factors, up to homotopy, through a cartesian diagram $E_n: \mathcal{S} \rightarrow \mathcal{D}$. The resulting sequence of morphisms

$$F \circ D \longrightarrow E_0 \longrightarrow T_\sigma(F) \circ D \longrightarrow E_1 \longrightarrow (T_\sigma)^2(F) \circ D \longrightarrow \dots$$

defines a sequential diagram (by Lemma D.9). Restricting along the inclusion $2\mathbb{N}_0 \subset \mathbb{N}_0$, we obtain a diagram where each morphism is homotopic to the corresponding one in diagram (2.3) and which is thus, by again Lemma D.9, equivalent to it. On the other hand, restricting along the inclusion $2\mathbb{N}_0 + 1 \subset \mathbb{N}_0$ yields a diagram E of the form

$$E_0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow \dots$$

whose colimit is cartesian by Lemma D.12, as each E_n is cartesian and \mathcal{D} is \mathcal{S} -differentiable. Since both inclusions are homotopy terminal, the colimit of the E_n is equivalent to the one of diagram (2.3), which finishes the proof. \square

Lemma 2.47. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, and \mathcal{D} an \mathcal{S} -differentiable ∞ -category. Then, for any functor $F: \mathcal{C} \rightarrow \mathcal{D}$, applying P_σ to $t_\sigma(F): F \rightarrow T_\sigma F$ yields an equivalence.*

Proof. Recall that $t_\sigma(F)$ is defined as the composition of the transformation $F \circ \tau_\sigma$, which is an equivalence as σ is full, and the upper horizontal map in the following homotopy commutative diagram

$$\begin{array}{ccc} \text{Res}_\emptyset \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} & \longrightarrow & \lim_{\mathcal{S}_{>\emptyset}} \circ \text{Res}_{\mathcal{S}_{>\emptyset}} \circ (F \circ) \circ \text{Lan}_\sigma \circ \text{Pad}_{\check{\mathcal{S}}} = T_\sigma(F) \\ \parallel & & \downarrow \simeq \\ F \circ (- \star \emptyset) & \longrightarrow & \lim_{s \in \mathcal{S}_{>\emptyset}} (F \circ (- \star s)) \end{array}$$

which we obtain from (the dual of) Lemma C.12. Hence it is enough to consider the lower horizontal map in the above diagram. Since, by Lemma 2.31, the functor P_σ preserves limits indexed by $\mathcal{S}_{>\emptyset}$, applying it to this map yields, by Lemma C.21, the upper triangle in the following homotopy commutative diagram in $\text{Fun}(\mathcal{C}, \mathcal{D})$

$$\begin{array}{ccc} & & P_\sigma(\lim_{s \in \mathcal{S}_{>\emptyset}} (F \circ (- \star s))) \\ & \nearrow & \downarrow \simeq \\ P_\sigma(F \circ (- \star \emptyset)) & \longrightarrow & \lim_{s \in \mathcal{S}_{>\emptyset}} P_\sigma(F \circ (- \star s)) \\ \simeq \downarrow & & \downarrow \simeq \\ P_\sigma(F) \circ (- \star \emptyset) & \longrightarrow & \lim_{s \in \mathcal{S}_{>\emptyset}} (P_\sigma(F) \circ (- \star s)) \end{array} \quad (2.4)$$

where the lower two vertical equivalences are provided by Lemma 2.31. This uses that $(- \star s)$ preserves left Kan extension along σ by Lemma C.16, for which we in turn use Lemma 2.22 to see that $(- \star s)$ preserves terminal objects and, for $s' \in \mathcal{S}$, colimits indexed by $\sigma \downarrow s'$ (since σ is full and hence $\sigma \downarrow s'$ has an initial object and is contractible).

But the lower horizontal map in diagram (2.4) is an equivalence since $P_\sigma(F)$ is σ -excisive by Lemma 2.46 and, for any $X \in \mathcal{C}$, the diagram $(X \star -) = (\text{Lan}_\sigma \text{Pad}_{\check{\mathcal{S}}})(X): \mathcal{S} \rightarrow \mathcal{C}$ is σ -cocartesian by Lemma C.1 (here we implicitly use that Res_X preserves limits and again Lemma C.21). This finishes the proof. \square

Lemma 2.48. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, and \mathcal{D} a \mathcal{S} -differentiable ∞ -category. Then, for any functor $F: \mathcal{C} \rightarrow \mathcal{D}$, applying P_σ to $p_\sigma(F): F \rightarrow P_\sigma F$ yields an equivalence.*

Proof. By Lemma 2.31, the functor P_σ preserves sequential colimits. Hence we have, by Lemma C.19, that the map $P_\sigma(p_\sigma(F))$ is an equivalence if and only if the structure map from $P_\sigma(F)$ to the colimit of the diagram

$$P_\sigma(F) \xrightarrow{P_\sigma(t_\sigma)} (P_\sigma T_\sigma)(F) \xrightarrow{P_\sigma(t_\sigma \circ T_\sigma)} (P_\sigma T_\sigma T_\sigma)(F) \longrightarrow \dots$$

is. But the latter follows from Lemma C.13 since each of the maps $P_\sigma(t_\sigma \circ (T_\sigma)^n)$ is an equivalence by Lemma 2.47. \square

Lemma 2.49. *Let C and D be objects of a 2-category, $l: C \rightarrow D$ and $r: D \rightarrow C$ morphisms, and $\eta: \text{id}_C \rightarrow r \circ l$ a 2-morphism. Assume that there is a 2-isomorphism*

$l \circ r \cong \text{id}_{\mathcal{D}}$, and that both $l \circ \eta$ and $\eta \circ r$ are 2-isomorphisms. Then there is an adjunction $l \dashv r$ with unit η .

Proof. This is [RV, Lemma B.4.2 and Remark B.4.3]. □

We are now ready to complete the proof of the main theorem:

Proof of Theorem 2.42. We want to apply Lemma 2.49 (in the homotopy 2-category of ∞ -categories) to $P_\sigma: \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Exc}_\sigma(\mathcal{C}, \mathcal{D})$ and the inclusion $\text{inc}: \text{Exc}_\sigma(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$. First note that P_σ actually lands in $\text{Exc}_\sigma(\mathcal{C}, \mathcal{D})$ by Lemma 2.46. Now, by Lemma 2.31, the transformation $p_\sigma: \text{id} \rightarrow \text{inc} \circ P_\sigma$ precomposed with inc is an equivalence. This also implies that $P_\sigma \circ \text{inc} \simeq \text{id}$ since $\text{Exc}_\sigma(\mathcal{C}, \mathcal{D})$ is a full subcategory. Furthermore the transformation $P_\sigma \circ p_\sigma$ is an equivalence by Lemma 2.48. □

3. Maps between approximations

As was mentioned in Remark 2.25, the construction of the universal σ -excisive approximation is not functorial in σ . However, by its universal property, we will get a map $P_\sigma(F) \rightarrow P_\tau(F)$ if $P_\tau(F)$ is σ -excisive. So, in this section, we will study what maps between (pre)shapes tell us about the relationship of the corresponding notions of excision.

One special case is the map $\lrcorner_n \rightarrow \lrcorner_{n+1}$ induced by the inclusion of $[n-1]$ into $[n]$. This will correspond to the classical fact that $(n-1)$ -excisive implies n -excisive (cf. [Goo92, Proposition 3.2] or [LurHA, Corollary 6.1.1.14]). The first subsection will focus on a generalization of this to more general (pre)shapes. However, under some conditions, a map $\sigma \rightarrow \tau$ of preshapes can also tell us that τ -excisive implies σ -excisive. This does not have an analogue in classical Goodwillie calculus (at least in the form we prove) and will be explored in the second subsection.

The names chosen for these concepts, indirect respectively direct maps, come from their effect on the universal excisive approximations. The mnemonic is that a direct map $\sigma \rightarrow \tau$ induces a map $P_\sigma \rightarrow P_\tau$ and the other way around for an indirect map.

Remark 3.1. In Remark 2.25, we saw that a map of (reduced) preshapes $(f, \check{f}): (\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}) \rightarrow (\tau: \check{\mathcal{T}} \rightarrow \mathcal{T})$ induces maps

$$T_\sigma(X) = \lim_{s \in \mathcal{S}_{>\emptyset}} (X \star_\sigma s) \longrightarrow \lim_{s \in \mathcal{S}_{>\emptyset}} (X \star_\tau f(s)) \longleftarrow \lim_{t \in \mathcal{T}_{>\emptyset}} (X \star_\tau t) = T_\tau(X)$$

which do not combine into a map between T_σ and T_τ . However, if one of these two maps were an equivalence for all X , then we would get a map in one direction. This can be seen as a motivation for there being two conditions, direct and indirect, one for each possible direction of the resulting map (and the conditions we give are basically chosen such that they guarantee one of the above maps to be an equivalence).

3.1. Indirect maps

Definition 3.2. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ and $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ be preshapes. A map $(f, \check{f}): \sigma \rightarrow \tau$ of preshapes is *indirect* if, for all $s \in \mathcal{S}$, the induced functor

$$\check{f}_s: \sigma \downarrow s \longrightarrow \tau \downarrow f(s), \quad \left(\check{s}, \sigma(\check{s}) \xrightarrow{l} s \right) \mapsto \left(\check{f}(\check{s}), \tau(\check{f}(\check{s})) = f(\sigma(\check{s})) \xrightarrow{f(l)} f(s) \right)$$

is homotopy terminal.

Lemma 3.3. Let $n \leq m$ be elements of \mathbb{N}_0 . Then the following is an indirect map of preshapes

$$\begin{array}{ccc} \mathcal{P}_{\leq 1}(n) & \xrightarrow{\check{f}} & \mathcal{P}_{\leq 1}(m) \\ \lrcorner_n \downarrow & & \downarrow \lrcorner_m \\ \mathcal{P}(n) & \xrightarrow{f} & \mathcal{P}(m) \end{array}$$

where f and \check{f} are induced by the inclusion of $[n-1]$ into $[m-1]$.

Proof. That it is a map of preshapes is clear. For indirectness we note that, for $S \in \mathcal{P}(n)$, the induced map $\downarrow_n S \rightarrow \downarrow_m f(S)$ is even an isomorphism. \square

Lemma 3.4. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ and $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ be preshapes, $(f, \check{f}): \sigma \rightarrow \tau$ an indirect map, and \mathcal{C} a σ -nice and τ -nice ∞ -category. Then, for any τ -cocartesian diagram $D: \mathcal{T} \rightarrow \mathcal{C}$, the diagram $\text{Res}_f(D)$ is σ -cocartesian.*

Proof. Consider the map (f, \check{f}) as the identity transformation in the following diagram

$$\begin{array}{ccc} \check{\mathcal{S}} & \xrightarrow{\check{f}} & \check{\mathcal{T}} \\ \sigma \downarrow & \text{id} \swarrow & \downarrow \tau \\ \mathcal{S} & \xrightarrow{f} & \mathcal{T} \end{array}$$

Applying $\text{Fun}(-, \mathcal{C})$ and taking the mate gives us a transformation $\text{id}_!: \text{Lan}_\sigma \text{Res}_{\check{f}} \rightarrow \text{Res}_f \text{Lan}_\tau$, which is an equivalence by Lemma C.6 and assumption. By Lemma A.2 we obtain a homotopy commutative diagram

$$\begin{array}{ccc} \text{Lan}_\sigma \text{Res}_{\check{f}} \text{Res}_\tau & \xrightarrow[\text{id}_!]{\simeq} & \text{Res}_f \text{Lan}_\tau \text{Res}_\tau \\ \parallel & & \downarrow \varepsilon_\tau \\ \text{Lan}_\sigma \text{Res}_\sigma \text{Res}_f & \xrightarrow{\varepsilon_\sigma} & \text{Res}_f \end{array}$$

where ε_σ and ε_τ are the counits of the respective adjunctions. Now note that ε_τ evaluated at D is an equivalence since D is τ -cocartesian. Hence ε_σ evaluated at $\text{Res}_f(D)$ is also an equivalence, which we wanted to show. \square

Proposition 3.5. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape, $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ a shape, \mathcal{C} a σ -nice and τ -nice ∞ -category, and \mathcal{D} an ∞ -category that admits limits indexed both by $\mathcal{S}_{>\emptyset}$ and by $\mathcal{T}_{>\emptyset}$. Furthermore, assume that there is an indirect map $(f, \check{f}): \sigma \rightarrow \tau$. Then each σ -excisive functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is also τ -excisive.*

Proof. Let $D: \mathcal{T} \rightarrow \mathcal{C}$ a τ -cocartesian diagram. We need to prove that $F \circ D$ is cartesian. For this we use the functor $p: \mathcal{T} \times \mathcal{S} \rightarrow \mathcal{T}$ given by $p(t, s) = t \amalg f(s)$ to transport diagrams indexed by \mathcal{T} to diagrams indexed by \mathcal{S} .

By Lemma 2.40 and τ being a shape, the diagram $D \circ (t \amalg -)$ is τ -cocartesian for any $t \in \mathcal{T}$. Then, by Lemma 3.4, the diagram $D \circ (t \amalg -) \circ f$ is σ -cocartesian. In particular this implies that $F \circ D \circ p: \mathcal{T} \times \mathcal{S} \rightarrow \mathcal{D}$ is cartesian when restricted to any $\{t\} \times \mathcal{S}$. Then, by Lemma D.13, we have that $F \circ D \circ p$ is already a limit diagram itself. Now, using that, by Lemma D.4, the restriction $p|_{(\mathcal{T} \times \mathcal{S})_{>\emptyset}}: (\mathcal{T} \times \mathcal{S})_{>\emptyset} \rightarrow \mathcal{T}_{>\emptyset}$ is homotopy initial, we obtain, by Lemma D.11, that $F \circ D$ is a limit diagram, which we wanted to show. \square

Together with Lemma 3.3 this implies the classical statement that n -excisive implies m -excisive for $n \leq m$:

Corollary 3.6. *Let $n \leq m$ be elements of \mathbb{N}_0 , \mathcal{C} an ∞ -category that admits all finite colimits, and \mathcal{D} an ∞ -category that admits limits indexed both by $\mathcal{P}(n)_{>\emptyset}$ and by $\mathcal{P}(m)_{>\emptyset}$. Then each $(n-1)$ -excisive functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is also $(m-1)$ -excisive.*

The following lemma will be needed later.

Lemma 3.7. *Let $\check{\mathcal{S}} \in \mathbf{Pos}$ and $\mathcal{S}, \mathcal{T} \in \mathbf{Pos}_{\Pi}$. Furthermore, let the following be a commutative diagram of functors between posets*

$$\begin{array}{ccc} & \check{\mathcal{S}} & \\ \sigma \swarrow & & \searrow \tau \\ \mathcal{S} & \xrightarrow{f} & \mathcal{T} \end{array}$$

such that τ is a shape and such that f is full and preserves finite coproducts. Then σ is also a shape that is finite if τ is, and $(f, \text{id}_{\check{\mathcal{S}}}): \sigma \rightarrow \tau$ is an indirect map of shapes.

Proof. First note that f is injective by Lemma D.1, hence \mathcal{S} is finite if \mathcal{T} is. Furthermore, this gives us that $\sigma^{-1}(\emptyset_{\mathcal{S}}) = \sigma^{-1}(f^{-1}(\emptyset_{\mathcal{T}})) = \tau^{-1}(\emptyset_{\mathcal{T}})$ is non-empty (note that $f^{-1}(\emptyset_{\mathcal{T}}) = \{\emptyset_{\mathcal{S}}\}$ as f is injective and preserves initial objects). In particular σ is a preshape.

We now need to show that, for all $s, t \in \mathcal{S}$ and $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \leq t \amalg s$, the full subposet

$$\check{\mathcal{S}}_{s,t,\check{k}}^{\sigma} = \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s \text{ and } \sigma(\check{k}) \leq t \amalg \sigma(\check{s})\} \subseteq \check{\mathcal{S}}$$

is contractible. For this note that our assumptions imply

$$\begin{aligned} \check{\mathcal{S}}_{s,t,\check{k}}^{\sigma} &= \{\check{s} \in \check{\mathcal{S}} \mid f(\sigma(\check{s})) \leq f(s) \text{ and } f(\sigma(\check{k})) \leq f(t \amalg \sigma(\check{s}))\} \\ &= \{\check{s} \in \check{\mathcal{S}} \mid \tau(\check{s}) \leq f(s) \text{ and } \tau(\check{k}) \leq f(t) \amalg \tau(\check{s})\} \\ &= \check{\mathcal{S}}_{f(s),f(t),\check{k}}^{\tau} \end{aligned}$$

which is contractible as τ is a shape and $\sigma(\check{k}) \leq t \amalg s$ implies $\tau(\check{k}) \leq f(t) \amalg f(s)$.

The tuple $(f, \text{id}_{\check{\mathcal{S}}})$ is a map of shapes since we have, as noted above, that $f^{-1}(\emptyset_{\mathcal{T}}) = \{\emptyset_{\mathcal{S}}\}$. For indirectness we need that, for all $s \in \mathcal{S}$, the functor $\sigma \downarrow s \rightarrow \tau \downarrow f(s)$ induced by $\text{id}_{\check{\mathcal{S}}}$ is homotopy terminal. We claim that it is even an isomorphism. For this, it is enough to show surjectivity since the functor in question is just the inclusion of one full subposet of $\check{\mathcal{S}}$ into another. This surjectivity is equivalent to $\tau(\check{s}) \leq f(s)$ implying $\sigma(\check{s}) \leq s$ for all $\check{s} \in \check{\mathcal{S}}$, which follows from the equality $f(\sigma(\check{s})) = \tau(\check{s})$ and f being full. \square

3.2. Direct maps

Definition 3.8. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ and $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ be preshapes. A map $(f, \check{f}): \sigma \rightarrow \tau$ of preshapes is *direct* if f is full and $f|_{\mathcal{S}_{>\emptyset}}: \mathcal{S}_{>\emptyset} \rightarrow \mathcal{T}_{>\emptyset}$ is homotopy initial.

Remark 3.9. Clearly any map of preshapes $(f, \check{f}): \sigma \rightarrow \tau$ such that f is an isomorphism is direct. Let us now furthermore assume that τ is full. In this case one can see very clearly

why τ -excisive should imply σ -excisive: any σ -cocartesian diagram is also τ -cocartesian. This is the case since, if we have a diagram $D: \mathcal{S} \rightarrow \mathcal{C}$ such that $D \simeq \text{Lan}_\sigma \text{Res}_\sigma D$, then also $D \simeq \text{Lan}_\tau \text{Lan}_{\check{f}} \text{Res}_\sigma D$, and thus D is τ -cocartesian by Lemma C.1. This can be seen as further motivation for the following proposition.

Proposition 3.10. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ and $\tau: \check{\mathcal{T}} \rightarrow \mathcal{T}$ be preshapes, \mathcal{C} a σ -nice and τ -nice ∞ -category, and \mathcal{D} an ∞ -category that admits limits indexed both by $\mathcal{S}_{>\emptyset}$ and by $\mathcal{T}_{>\emptyset}$. Assume that τ is a shape or full and that there is a direct map $(f, \check{f}): \sigma \rightarrow \tau$. Then each τ -excisive functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is also σ -excisive.*

Proof. Let $D: \mathcal{S} \rightarrow \mathcal{C}$ be a σ -cocartesian diagram. We need to show that $F \circ D$ is cartesian. For this we use Lan_f to transport D to a diagram indexed by \mathcal{T} .

We have

$$\text{Lan}_f(D) \simeq (\text{Lan}_f \text{Lan}_\sigma \text{Res}_\sigma)(D) \simeq (\text{Lan}_\tau \text{Lan}_{\check{f}} \text{Res}_\sigma)(D)$$

which is τ -cocartesian. This follows from Lemma C.1 if τ is full and from Lemma 2.40 if τ is a shape. In particular, we obtain that $F \circ \text{Lan}_f(D)$ is cartesian. Hence, by Lemma D.11, the restriction $F \circ \text{Lan}_f(D) \circ f$ is cartesian (here we use that $f|_{\mathcal{S}_{>\emptyset}}$ is homotopy initial). Now, since f is full and hence fully faithful, we have that

$$F \circ \text{Lan}_f(D) \circ f = F \circ (\text{Res}_f \text{Lan}_f)(D) \simeq F \circ D$$

and thus that $F \circ D$ is cartesian, which we wanted to show. \square

Remark 3.11. The condition that f is full is probably a bit stronger than actually required. We do not need that $\text{Res}_f \text{Lan}_f D \simeq D$ for all $D: \mathcal{S} \rightarrow \mathcal{C}$, but only for those that are σ -cocartesian. Possibly a weaker condition guaranteeing this could be formulated.

3.3. Example application: punctured cubes

Our setup allows us to study more families of (pre)shapes than just the usual inclusion of cubes \sqcup_n . One such family arises when, instead of the \sqcup_n , we consider the inclusions of a punctured cube (i.e. without the terminal object) into the whole cube. These have some relevance, for example, by (the dual of) Remark 2.3 a diagram is cocartesian with respect to this inclusion if and only if it is cocartesian in the classical sense (i.e the one dual to Definition 2.2). In this subsection we will exemplarily show that the corresponding excision condition for a larger cube implies the one for a smaller cube, as it nicely uses both direct and indirect maps.

Notation 3.12. Let $n \in \mathbb{N}_{\geq 1}$. We denote by $\mathcal{P}_{<^*}(n)$ the full subposet of $\mathcal{P}(n)$ consisting of the non-terminal objects and by $\square_n: \mathcal{P}_{<^*}(n) \rightarrow \mathcal{P}(n)$ the inclusion.

Remark 3.13. It follows directly from Example 2.37 that \square_n is a shape for any $n \in \mathbb{N}_{\geq 1}$.

Remark 3.14. Note that \square_2 is isomorphic to \sqcup_2 .

Proposition 3.15. *Let $n \geq m$ be elements of $\mathbb{N}_{\geq 1}$, \mathcal{C} an ∞ -category that admits all finite colimits, and \mathcal{D} an ∞ -category that admits limits indexed both by $\mathcal{P}(n)_{>\emptyset}$ and by $\mathcal{P}(m)_{>\emptyset}$. Then each \square_n -excisive functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is also \square_m -excisive.*

Proof. Let $s: [n-1] \rightarrow [m-1]$ be a surjective map. By taking images this induces a functor $\mathcal{P}(n) \rightarrow \mathcal{P}(m)$ which we also denote by s . We write $\mathcal{Q} := s^{-1}(\mathcal{P}_{<*(m)})$ considered as a full subposet of $\mathcal{P}(n)$ and note that it is clear that the inclusion $\iota: \mathcal{Q} \rightarrow \mathcal{P}(n)$ is a preshape.

Claim 3.15.1. *The map of preshapes $(s, \check{s}): \iota \rightarrow \square_m$ is indirect, where $\check{s}: \mathcal{Q} \rightarrow \mathcal{P}_{<*(m)}$ is the restriction of s .*

Proof. We need to show that for all $N \subseteq [n-1]$ the induced map $\check{s}_N: \iota \downarrow N \rightarrow \square_m \downarrow s(N)$ is homotopy terminal, i.e. that for all $M \in \square_m \downarrow s(N)$ the category

$$M \downarrow \check{s}_N = \{N' \subseteq [n-1] \mid N' \subseteq N \text{ and } M \subseteq s(N') \subsetneq [m-1]\} \subseteq \mathcal{Q}$$

is contractible. For this purpose, define

$$\mathcal{K} := \{K \subsetneq [m-1] \mid M \subseteq K\} \subseteq \mathcal{P}_{<*(m)}$$

considered as a full subposet. Note that \mathcal{K} has the initial object M , hence \mathcal{K} is contractible, and it is enough to show that it is homotopy equivalent to $M \downarrow \check{s}_N$. For this, let $l: M \downarrow \check{s}_N \rightarrow \mathcal{K}$ be given by s and $r: \mathcal{K} \rightarrow M \downarrow \check{s}_N$ by $K \mapsto s^{-1}(K) \cap N$. To see that r is well-defined note that $s(s^{-1}(K) \cap N) \subseteq K \subsetneq [m-1]$ and that, since $M \subseteq s(N)$, there exists for each $a \in M$ an element $b \in N$ such that $s(b) = a \in K$, hence $b \in s^{-1}(K)$ and $a \in s(s^{-1}(K) \cap N)$. We claim that l is left adjoint to r . For this we need to show that, for all $N' \in M \downarrow \check{s}_N$ and $K \in \mathcal{K}$, we have $s(N') \subseteq K$ if and only if $N' \subseteq s^{-1}(K) \cap N$, which is clear. \blacksquare

By Proposition 3.5, we now know that a ι -excisive functor $\mathcal{C} \rightarrow \mathcal{D}$ is \square_m -excisive. But the map of preshapes $(\text{id}, \check{\iota}): \iota \rightarrow \square_n$ is direct, where $\check{\iota}: \mathcal{Q} \rightarrow \mathcal{P}_{<*(n)}$ is the restriction of ι . Hence, by Proposition 3.10, a \square_n -excisive functor $\mathcal{C} \rightarrow \mathcal{D}$ is ι -excisive, and we are done. \square

Note, in particular, that \square_n -excisive implies \square_2 -excisive and hence 1-excisive for any $n \in \mathbb{N}_{\geq 2}$. This implication can be turned around when one only considers functors that additionally preserve terminal objects (under some pointedness conditions). The following proof of this fact was worked out together with Tashi Walde (though any mistake is mine).

Proposition 3.16. *Let \mathcal{C} be an ∞ -category that admits all finite colimits and is pointed (i.e. it has an object that is both initial and terminal), \mathcal{D} a differentiable pointed ∞ -category, $F: \mathcal{C} \rightarrow \mathcal{D}$ a 1-homogeneous functor (i.e. it is 1-excisive and preserves terminal objects), and $n \in \mathbb{N}_{\geq 1}$. Then F is \square_n -excisive.*

Proof. First assume that \mathcal{D} is stable (cf. [LurHA, Definition 1.1.1.9]). By Remark 2.3, a functor is \square_n -excisive if and only if it sends cocartesian n -cubes (in the sense dual to Definition 2.2) to cartesian ones. But, by [LurHA, Proposition 1.2.4.13], in a stable ∞ -category an n -cube is cartesian if and only if it is cocartesian. In particular the 1-excisive functor F preserves pushouts. Since it also preserves initial objects (as \mathcal{C} is pointed), it already preserves all finite colimits by [LurHTT, Corollary 4.4.2.5]. Hence F sends cocartesian n -cubes in \mathcal{C} to cocartesian n -cubes in \mathcal{D} , the latter of which are also cartesian. (Note that in this case, i.e. when \mathcal{D} is stable, we do not need the differentiability condition as we only used that \mathcal{D} admits all finite limits and colimits, which is automatic by [LurHA, Proposition 1.1.3.4].)

For the general case, we use that, by [LurHA, Corollary 6.1.2.9], the 1-homogeneous functor F factors (up to equivalence) as a composition

$$\mathcal{C} \xrightarrow{F'} \mathrm{Sp}(\mathcal{D}) \xrightarrow{\Omega^\infty} \mathcal{D}$$

where $\mathrm{Sp}(\mathcal{D})$ is the ∞ -category of spectrum objects in \mathcal{D} (cf. [LurHA, Definition 1.4.2.8]), Ω^∞ is as in [LurHA, Notation 1.4.2.20], and F' is a 1-homogeneous functor. Since $\mathrm{Sp}(\mathcal{D})$ is stable by [LurHA, Corollary 1.2.4.17] we obtain by the above special case that F' is \square_n -excisive. But Ω^∞ preserves finite limits since it is given by an evaluation (here we implicitly use [LurHA, Remark 1.4.2.3]). Thus the composition $\Omega^\infty \circ F' \simeq F$ is also \square_n -excisive. \square

Remark 3.17. It would actually be enough to assume \mathcal{C} to just have a terminal object instead of being pointed. To see this one would need to show that if a functor preserves pushouts, then it already preserves colimits indexed by $\mathcal{P}_{<*(n)}$ for all $n \in \mathbb{N}_{\geq 1}$. This is not too hard, but we omit the rather technical proof.

4. The structure of shapes

4.1. Full shapes

Notation 4.1. Let $f: \mathcal{P} \rightarrow \mathcal{Q}$ be a map of posets. We denote by $\text{im } f \subseteq \mathcal{Q}$ the essential image of f , i.e. the poset with elements $f(\mathcal{P})$ and partial order \preceq generated by the relation $q \preceq q'$ if there are $p \in f^{-1}(q)$ and $p' \in f^{-1}(q')$ such that $p \leq p'$.

Remark 4.2. Note that $\text{im } f$ is a subposet of \mathcal{Q} but does not need to be full. To distinguish the different partial orders in this situation we will use, as in the definition, the symbol \preceq for the one of $\text{im } f$.

Definition 4.3. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape. We write $\tilde{\sigma}: \text{im } \sigma \rightarrow \mathcal{S}$ for the preshape occurring in the factorization

$$\begin{array}{ccc} & \text{im } \sigma & \\ \nearrow & & \searrow \tilde{\sigma} \\ \check{\mathcal{S}} & \xrightarrow{\sigma} & \mathcal{S} \end{array}$$

Remark 4.4. That $\tilde{\sigma}$ is again a preshape is immediate as $\emptyset_{\mathcal{S}} \in \text{im } \sigma$.

Lemma 4.5. *If $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ is a finite shape, then $\tilde{\sigma}: \text{im } \sigma \rightarrow \mathcal{S}$ is full.*

Proof. We need to show that for any two elements i and j of $\text{im } \sigma$ such that $i \leq j$, we also have $i \preceq j$. We do this by induction on

$$d(i, j) := \sup\{n \in \mathbb{N}_0 \mid \exists s_0, \dots, s_n \in \mathcal{S} \text{ such that } i = s_0 \preceq \dots \preceq s_n = j\}$$

which is finite since \mathcal{S} is finite by assumption.

If $d(i, j) = 0$, then $i = j$ and we are done. Otherwise, let $\check{k} \in \sigma^{-1}(i)$. Then, by the definition of a shape, the full subposet

$$\check{\mathcal{S}}_{j, \emptyset, \check{k}} = \{\check{i} \in \check{\mathcal{S}} \mid i \leq \sigma(\check{i}) \leq j\} \subseteq \check{\mathcal{S}}$$

is contractible, in particular connected, as $\sigma(\check{k}) = i \leq j = j \amalg \emptyset$. Since $i \neq j$, the preimage $\sigma^{-1}(i)$ is a proper non-empty subset of $\check{\mathcal{S}}_{j, \emptyset, \check{k}}$. Thus, there must be a morphism in $\check{\mathcal{S}}_{j, \emptyset, \check{k}}$ with exactly one of target or source lying in $\sigma^{-1}(i)$. But if it were the target, there would be an element $\check{i} \in \check{\mathcal{S}}_{j, \emptyset, \check{k}}$ such that $\sigma(\check{i}) \preceq i$, a contradiction. So there must be $\check{i} \in \sigma^{-1}(i)$ and $\check{l} \in \check{\mathcal{S}}_{j, \emptyset, \check{k}} \setminus \sigma^{-1}(i)$ such that $\check{i} \leq \check{l}$. In particular $i = \sigma(\check{i}) \preceq \sigma(\check{l}) \leq j$, hence it is enough to show that $\sigma(\check{l}) \preceq j$. But as $i \neq \sigma(\check{l})$ we have $d(\sigma(\check{l}), j) \preceq d(i, j)$, so the statement follows by induction. \square

Lemma 4.6. *A finite shape $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ is full if and only if it is injective.*

Proof. The “only if” part follows from Lemma D.1. For the other direction note that, if σ is injective, then $\sigma: \check{\mathcal{S}} \rightarrow \text{im } \sigma$ is an isomorphism since the generating relation in Definition 4.3 is, in this case, already transitive. Hence, by the factorization

$$\begin{array}{ccc} & \text{im } \sigma & \\ \cong \nearrow & & \searrow \tilde{\sigma} \\ \check{\mathcal{S}} & \xrightarrow{\sigma} & \mathcal{S} \end{array}$$

and $\tilde{\sigma}$ being full by Lemma 4.5, we obtain that σ is full as well. \square

The following result tells us that for each finite shape there is a full finite shape with the same excision properties.

Proposition 4.7. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a finite shape. Then $\tilde{\sigma}: \text{im } \sigma \rightarrow \mathcal{S}$ is again a finite shape (and full, by Lemma 4.5) and the map of shapes*

$$\begin{array}{ccc} \check{\mathcal{S}} & \xrightarrow{\sigma} & \text{im } \sigma \\ \sigma \downarrow & & \downarrow \tilde{\sigma} \\ \mathcal{S} & \xrightarrow{\text{id}} & \mathcal{S} \end{array} \quad (4.1)$$

is both direct and indirect. In particular, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories is σ -excisive if and only if it is $\tilde{\sigma}$ -excisive (as long as \mathcal{C} is σ -nice (which implies $\tilde{\sigma}$ -nice), and \mathcal{D} admits limits indexed by $\mathcal{S}_{>\emptyset}$).

Proof. For the first part we need to show that, for all $s, t \in \mathcal{S}$ and $\tilde{k} \in \text{im } \sigma$ such that $\tilde{\sigma}(\tilde{k}) \leq t \amalg s$, the full subposet

$$\tilde{\mathcal{I}}_{s,t,\tilde{k}} := \{\tilde{i} \in \text{im } \sigma \mid \tilde{\sigma}(\tilde{i}) \leq s \text{ and } \tilde{\sigma}(\tilde{k}) \leq t \amalg \tilde{\sigma}(\tilde{i})\} \subseteq \text{im } \sigma$$

is contractible. Let $\check{l} \in \check{\sigma}^{-1}(\tilde{k})$. Then, as σ is a shape and $\sigma(\check{l}) \leq t \amalg s$, we know that the full subposet

$$\check{\mathcal{S}}_{s,t,\check{l}} := \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s \text{ and } \tilde{\sigma}(\tilde{k}) = \sigma(\check{l}) \leq t \amalg \sigma(\check{s})\} \subseteq \check{\mathcal{S}}$$

is contractible. By commutativity of diagram (4.1), we have that $\check{\sigma}$ induces a functor $g: \check{\mathcal{S}}_{s,t,\check{l}} \rightarrow \tilde{\mathcal{I}}_{s,t,\tilde{k}}$. It is now enough to check that g is a homotopy equivalence, in particular that it is homotopy initial, i.e. that for each $\tilde{i} \in \tilde{\mathcal{I}}_{s,t,\tilde{k}}$ the full subposet

$$\mathcal{G}_{\tilde{i}} := g \downarrow \tilde{i} = \{\check{s} \in \check{\mathcal{S}}_{s,t,\check{l}} \mid g(\check{s}) \preceq \tilde{i}\} \subseteq \check{\mathcal{S}}_{s,t,\check{l}}$$

is contractible. By Lemma 4.5, the condition $g(\check{s}) \preceq \tilde{i}$ is equivalent to $\sigma(\check{s}) = \tilde{\sigma}(g(\check{s})) \leq \tilde{\sigma}(\tilde{i})$. Hence, using that $\tilde{\sigma}(\tilde{i}) \leq s$, we obtain $\mathcal{G}_{\tilde{i}} = \check{\mathcal{S}}_{\tilde{\sigma}(\tilde{i}),t,\check{l}}$, which is contractible since σ is a shape and $\sigma(\check{l}) = \tilde{\sigma}(\tilde{k}) \leq t \amalg \tilde{\sigma}(\tilde{i})$ by definition of $\tilde{\mathcal{I}}_{s,t,\tilde{k}}$.

That $(\text{id}, \tilde{\sigma})$ is direct is clear. For indirectness we have to show that for any $s \in \mathcal{S}$ the functor $\tilde{\sigma}_s: \sigma \downarrow s \rightarrow \tilde{\sigma} \downarrow s$ is homotopy terminal, i.e. that for any $\tilde{s} \in \tilde{\sigma} \downarrow s$ the category $\tilde{s} \downarrow \tilde{\sigma}_s$ is contractible. This category $\tilde{s} \downarrow \tilde{\sigma}_s$ can be identified with the full subposet

$$\mathcal{H}_{s, \tilde{s}} = \{\check{t} \in \check{\mathcal{S}} \mid \sigma(\check{t}) \leq s \text{ and } \tilde{s} \preceq \check{\sigma}(\check{t})\} \subseteq \check{\mathcal{S}}.$$

Let $\check{k} \in \check{\sigma}^{-1}(\tilde{s})$. Then we have, again using Lemma 4.5, that $\mathcal{H}_{s, \tilde{s}}$ is just $\check{\mathcal{S}}_{s, \emptyset, \check{k}}$ and hence contractible since σ is a shape and $\sigma(\check{k}) = \tilde{\sigma}(\tilde{s}) \leq s = \emptyset \amalg s$ by definition of \tilde{s} . \square

As a corollary we obtain the following version of Theorem 2.42 for shapes which are finite but not necessarily full.

Corollary 4.8. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a finite shape, \mathcal{C} a σ -nice ∞ -category with a terminal object, and \mathcal{D} an \mathcal{S} -differentiable ∞ -category. Then there is an adjunction with left adjoint $P_{\tilde{\sigma}}: \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Exc}_{\tilde{\sigma}}(\mathcal{C}, \mathcal{D}) = \text{Exc}_{\sigma}(\mathcal{C}, \mathcal{D})$, right adjoint the inclusion $\text{inc}: \text{Exc}_{\sigma}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$, and unit $p_{\tilde{\sigma}}: \text{id} \rightarrow \text{inc} \circ P_{\tilde{\sigma}}$.*

4.2. Free shapes

Definition 4.9. Let \mathcal{P} be a poset and $M \subseteq \mathcal{P}$ a subset. We write

- a) $\downarrow M := \{x \in \mathcal{P} \mid \exists y \in M: x \leq y\}$ for the *down-set* of M .
- b) $\mathcal{O}(\mathcal{P}) := \{N \subseteq \mathcal{P} \mid N = \downarrow N\}$ for the *down-set lattice*, i.e. the poset of downward closed subsets ordered by inclusion.
- c) $o_{\mathcal{P}}: \mathcal{P} \rightarrow \mathcal{O}(\mathcal{P})$ for the canonical functor given by $x \mapsto \downarrow\{x\}$.

It is well-known (c.f. [DP, Examples 2.6 (3)]) that $\mathcal{O}(\mathcal{P})$ is a complete lattice, in particular that it has all coproducts, and that these coproducts are given by taking the union of subsets. Actually, the map $o_{\mathcal{P}}$ is even the universal map from \mathcal{P} to a poset with all coproducts.

The following definition introduces a similar construction which adds coproducts to posets that already have initial objects. Just taking \mathcal{O} would add a new initial object (the empty set) which we do not want.

Notation 4.10. Let $\mathcal{P} \in \mathbf{Pos}_{\emptyset}$. We write $\tilde{\mathcal{O}}(\mathcal{P}) \subsetneq \mathcal{O}(\mathcal{P})$ for the full subposet of non-empty subsets. Noting that $o_{\mathcal{P}}$ uniquely factors as $\mathcal{P} \rightarrow \tilde{\mathcal{O}}(\mathcal{P}) \subsetneq \mathcal{O}(\mathcal{P})$, we will also, by abuse of notation, write $o_{\mathcal{P}}: \mathcal{P} \rightarrow \tilde{\mathcal{O}}(\mathcal{P})$ for the first map in this factorization.

Remark 4.11. Note that $\tilde{\mathcal{O}}(\mathcal{P})$ still has all coproducts: non-empty ones are again given by taking unions of subsets and the initial object $\{\emptyset_{\mathcal{P}}\}$ constitutes an empty one.

Lemma 4.12. *For $\mathcal{P} = \mathcal{P}_{\leq 1}(n)$, there is a canonical isomorphism of posets under $\mathcal{P}_{\leq 1}(n)$ between $o_{\mathcal{P}_{\leq 1}(n)}: \mathcal{P}_{\leq 1}(n) \rightarrow \tilde{\mathcal{O}}(\mathcal{P}_{\leq 1}(n))$ and $\perp_n: \mathcal{P}_{\leq 1}(n) \rightarrow \mathcal{P}(n)$.*

Proof. This follows directly from the definitions. \square

The following lemma now makes precise what we claimed before about the universality of this construction.

Lemma 4.13. *The construction $\tilde{\mathcal{O}}$ extends to a functor $\mathbf{Pos}_{\emptyset} \rightarrow \mathbf{Pos}_{\mathbb{I}}$ by defining, for $f \in \mathbf{Hom}_{\mathbf{Pos}_{\emptyset}}(\mathcal{P}, \mathcal{P}')$, the induced map $\tilde{\mathcal{O}}(f)$ to be given by $M \mapsto \downarrow f(M)$.*

Furthermore $\tilde{\mathcal{O}}$ is left adjoint to the forgetful functor $U: \mathbf{Pos}_{\mathbb{I}} \rightarrow \mathbf{Pos}_{\emptyset}$. More explicitly, for all posets $\mathcal{P} \in \mathbf{Pos}_{\emptyset}$, $\mathcal{Q} \in \mathbf{Pos}_{\mathbb{I}}$, and any $f \in \mathbf{Hom}_{\mathbf{Pos}_{\emptyset}}(\mathcal{P}, U(\mathcal{Q}))$, the functor $u_f: \tilde{\mathcal{O}}(\mathcal{P}) \rightarrow \mathcal{Q}$ given by $M \mapsto \coprod_{m \in M} f(m)$ is the unique element of $\mathbf{Hom}_{\mathbf{Pos}_{\mathbb{I}}}(\tilde{\mathcal{O}}(\mathcal{P}), \mathcal{Q})$ such that

$$\begin{array}{ccc} & \mathcal{P} & \\ o_{\mathcal{P}} \swarrow & & \searrow f \\ \tilde{\mathcal{O}}(\mathcal{P}) & \xrightarrow{\quad} & \mathcal{Q} \end{array}$$

commutes, and the assignment $f \mapsto u_f$ is natural in both \mathcal{P} and \mathcal{Q} .

Moreover o is a natural transformation $\text{id} \rightarrow U \circ \tilde{\mathcal{O}}$ of functors $\mathbf{Pos}_{\emptyset} \rightarrow \mathbf{Pos}_{\emptyset}$.

Proof. First note that it is clear that $\tilde{\mathcal{O}}(f)$ is a morphism in $\mathbf{Pos}_{\mathbb{I}}$. For functoriality of $\tilde{\mathcal{O}}$ we furthermore need that, for $M \in \tilde{\mathcal{O}}(\mathcal{P})$ and $f: \mathcal{P} \rightarrow \mathcal{Q}$ and $g: \mathcal{Q} \rightarrow \mathcal{R}$ maps of posets, we have $M = \downarrow \text{id}(M)$ and $\downarrow(g \circ f)(M) = \downarrow g(\downarrow f(M))$. The former statement, as well as the inclusion \subseteq of the latter, are clear. For the other inclusion let $x \in \mathcal{R}$ and note that $x \leq g(y)$ for some $y \in \downarrow f(M)$ implies that there is $z \in M$ such that $y \leq f(z)$. Hence $x \leq g(f(z))$ and thus $x \in \downarrow g(f(M))$.

For naturality of o we need that, for $x \in \mathcal{P}$ and $f \in \mathbf{Hom}_{\mathbf{Pos}_{\emptyset}}(\mathcal{P}, \mathcal{Q})$, we have $\downarrow f(\downarrow \{x\}) = \downarrow \{f(x)\}$, which follows from the same argument as the functoriality.

For the adjunction, it is clear that u_f is a functor and that $u_f \circ o_{\mathcal{P}} = f$ as we have, for $x \in \mathcal{P}$, that $\coprod_{y \leq x} f(y) = f(x)$ (this uses that in a poset the coproduct of a set of objects with a maximum is that maximum). To see that u_f is a morphism in $\mathbf{Pos}_{\mathbb{I}}$, note that, for any non-empty subset $L \subseteq \tilde{\mathcal{O}}(\mathcal{P})$, we have

$$u_f \left(\coprod_{M \in L} M \right) = u_f \left(\bigcup_{M \in L} M \right) = \coprod_{m \in \bigcup_{M \in L} M} m = \coprod_{M \in L} \coprod_{m \in M} m = \coprod_{M \in L} u_f(M),$$

where the third equation follows from the fact that, in a poset, a coproduct over multiple copies of the same object is that object again. For the empty coproduct we have $u_f(\{\emptyset_{\mathcal{P}}\}) = f(\emptyset_{\mathcal{P}}) = \emptyset_{\mathcal{Q}}$ as f preserves initial objects by assumption. For uniqueness, note that any element of $\tilde{\mathcal{O}}(\mathcal{P})$ is the coproduct of elements in the image of $o_{\mathcal{P}}$. Hence any element of $\mathbf{Hom}_{\mathbf{Pos}_{\mathbb{I}}}(\tilde{\mathcal{O}}(\mathcal{P}), \mathcal{Q})$ is already uniquely determined by its restriction along $o_{\mathcal{P}}$.

For naturality of the map $f \mapsto u_f$ in the variable \mathcal{P} we need that $u_{f \circ p} = u_f \circ \tilde{\mathcal{O}}(p)$ for all morphisms p in \mathbf{Pos}_{\emptyset} , which follows from $\coprod_{m \in M} f(p(m)) = \coprod_{m' \in \downarrow p(M)} f(m')$. For naturality in the variable \mathcal{Q} we need that $u_{q \circ f} = q \circ u_f$ for all morphisms q in $\mathbf{Pos}_{\mathbb{I}}$, which follows from q preserving coproducts. \square

Lemma 4.14. *For any $\check{\mathcal{S}} \in \mathbf{Pos}_{\emptyset}$ the map $o_{\check{\mathcal{S}}}: \check{\mathcal{S}} \rightarrow \tilde{\mathcal{O}}(\check{\mathcal{S}})$ is a full shape. Moreover, if $\check{\mathcal{S}}$ is finite, then $o_{\check{\mathcal{S}}}$ is also finite.*

Proof. First note that it is clear that $o_{\check{\mathcal{S}}}$ is a preshape and that $\tilde{\mathcal{O}}(\check{\mathcal{S}})$ is finite if $\check{\mathcal{S}}$ is. Fullness follows from the fact that, for \check{a} and \check{b} in $\check{\mathcal{S}}$, the inclusion $\downarrow\{\check{a}\} \subseteq \downarrow\{\check{b}\}$ implies $\check{a} \leq \check{b}$.

Now we want to employ Lemma 2.36 to show that $o_{\check{\mathcal{S}}}$ is a shape. For this we need that, for all $A, B \in \tilde{\mathcal{O}}(\check{\mathcal{S}})$ and $\check{c} \in \check{\mathcal{S}}$ such that $o_{\check{\mathcal{S}}}(\check{c}) \leq A \amalg B$, we have $o_{\check{\mathcal{S}}}(\check{c}) \leq A$ or $o_{\check{\mathcal{S}}}(\check{c}) \leq B$. But $\downarrow\{\check{c}\} \subseteq A \cup B$ implies $\check{c} \in A$ or $\check{c} \in B$. In both cases the set $\downarrow\{\check{c}\}$ must already lie in one of A and B as they are downward closed. This implies the statement. \square

Remark 4.15. Combining Lemma 4.14 with Lemma 4.12, we obtain as a special case the fact that $\lrcorner_n: \mathcal{P}_{\leq 1}(n) \rightarrow \mathcal{P}(n)$ is a full shape (which we had already seen in Example 2.37).

The following result tells us in particular that, for any finite poset $\check{\mathcal{S}}$ with more than one object, there exists $n \in \mathbb{N}_0$ such that $o_{\check{\mathcal{S}}}$ -excisive is equivalent to n -excisive.

Proposition 4.16. *Let $\check{\mathcal{S}} \in \mathbf{Pos}_{\emptyset}$ and $\mathcal{M} \subseteq \check{\mathcal{S}}$ a non-empty full downward closed subposet such that, for all $\check{s} \in \check{\mathcal{S}}_{>\emptyset}$, there exists an element $m \in \mathcal{M}_{>\emptyset}$ with $m \leq \check{s}$ (if $\check{\mathcal{S}}$ is finite this is equivalent to requiring \mathcal{M} to contain all minimal elements of $\check{\mathcal{S}}_{>\emptyset}$). Then the inclusion $i: \mathcal{M} \rightarrow \check{\mathcal{S}}$ induces a map of shapes*

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{i} & \check{\mathcal{S}} \\ o_{\mathcal{M}} \downarrow & & \downarrow o_{\check{\mathcal{S}}} \\ \tilde{\mathcal{O}}(\mathcal{M}) & \xrightarrow{\tilde{\mathcal{O}}(i)} & \tilde{\mathcal{O}}(\check{\mathcal{S}}) \end{array}$$

which is both direct and indirect.

In particular, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories is $o_{\check{\mathcal{S}}}$ -excisive if and only if it is $o_{\mathcal{M}}$ -excisive (as long as \mathcal{C} is $o_{\check{\mathcal{S}}}$ -nice (which implies $o_{\mathcal{M}}$ -nice), and \mathcal{D} admits limits indexed by both $\tilde{\mathcal{O}}(\mathcal{M})_{>\emptyset}$ and $\tilde{\mathcal{O}}(\check{\mathcal{S}})_{>\emptyset}$).

Proof. First note that the square commutes by naturality of o .

Furthermore $\tilde{\mathcal{O}}(i)$ is full since we have, for any $M \in \tilde{\mathcal{O}}(\mathcal{M})$, that $\downarrow i(M) = i(M)$ (as \mathcal{M} is downward closed) and hence that $\tilde{\mathcal{O}}(i)(M) \subseteq \tilde{\mathcal{O}}(i)(N)$ implies $M \subseteq N$. Thus, for directness, it only remains to show that $\tilde{\mathcal{O}}(i)_{>\emptyset}: \tilde{\mathcal{O}}(\mathcal{M})_{>\emptyset} \rightarrow \tilde{\mathcal{O}}(\check{\mathcal{S}})_{>\emptyset}$ is homotopy initial. For this let $L \in \tilde{\mathcal{O}}(\check{\mathcal{S}})_{>\emptyset}$. Then $\tilde{\mathcal{O}}(i)_{>\emptyset} \downarrow L$ has the terminal object $\{m \in \mathcal{M} \mid i(m) \in L\}$ (which is actually contained in $\tilde{\mathcal{O}}(\mathcal{M})_{>\emptyset}$ since our assumptions imply that there exists an element $m \in \mathcal{M}_{>\emptyset}$ with $i(m) \in L$) and is thus contractible.

For indirectness we have to check that, for all $M \in \tilde{\mathcal{O}}(\mathcal{M})$, the functor $i_M: o_{\mathcal{M}} \downarrow M \rightarrow o_{\check{\mathcal{S}}} \downarrow \tilde{\mathcal{O}}(i)(M)$ induced by i is homotopy terminal. But, by \mathcal{M} being downward closed in $\check{\mathcal{S}}$, we have that $\downarrow\{\check{s}\} \subseteq \downarrow i(M)$ implies $\check{s} \in M$. Hence i_M is even an isomorphism. \square

Corollary 4.17. *Let $\check{\mathcal{S}} \in \mathbf{Pos}_{\emptyset}$ be finite. Then a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories is $o_{\check{\mathcal{S}}}$ -excisive if and only if it is $(n-1)$ -excisive, where n is the cardinality of the discrete poset \mathcal{M} of minimal elements of $\check{\mathcal{S}}_{>\emptyset}$ (as long as \mathcal{C} admits all finite colimits, and \mathcal{D} admits limits indexed by both $\mathcal{P}(n)_{>\emptyset}$ and $\tilde{\mathcal{O}}(\check{\mathcal{S}})_{>\emptyset}$).*

Proof. This follows from Proposition 4.16 applied to $\mathcal{M} \cup \{\emptyset\}$ by noting that $\mathcal{M} \cup \{\emptyset\} \cong \mathcal{P}_{\leq 1}(n)$ and using Lemma 4.12. \square

4.3. Boring shapes

Definition 4.18. Let $\mathcal{P} \in \mathbf{Pos}_\emptyset$, $\mathcal{Q} \in \mathbf{Pos}_\Pi$, and $f \in \mathbf{Hom}_{\mathbf{Pos}_\emptyset}(\mathcal{P}, \mathcal{Q})$. We write $v_f: \mathcal{P} \rightarrow \text{im } u_f$ and $w_f: \text{im } u_f \rightarrow \mathcal{Q}$ for the maps occurring in the following factorizations

$$\begin{array}{ccc}
 & \mathcal{P} & \\
 \swarrow \sigma_{\mathcal{P}} & \downarrow v_f & \searrow f \\
 \tilde{\mathcal{O}}(\mathcal{P}) & \xrightarrow{u_f} & \mathcal{Q} \\
 \searrow & \downarrow & \swarrow w_f \\
 & \text{im } u_f &
 \end{array}$$

of f .

This can not be applied to $f = \sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ an arbitrary (pre)shape as $\check{\mathcal{S}}$ does not necessarily have an initial object, so we need to restrict ourselves to reduced (pre)shapes. Luckily, by Proposition 4.7, we do not lose much by doing so (at least in the finite case).

Lemma 4.19. Let $\mathcal{P} \in \mathbf{Pos}_\emptyset$, $\mathcal{Q} \in \mathbf{Pos}_\Pi$, and $f \in \mathbf{Hom}_{\mathbf{Pos}_\emptyset}(\mathcal{P}, \mathcal{Q})$. Then $\text{im } u_f$ has all coproducts and $w_f: \text{im } u_f \rightarrow \mathcal{Q}$ is full and preserves coproducts.

Proof. We first show fullness. To this end, let M and M' be elements of $\tilde{\mathcal{O}}(\mathcal{P})$ such that $u_f(M) \leq u_f(M')$. But then, since u_f preserves coproducts, we have $u_f(M) \preceq u_f(M \cup M') = u_f(M) \amalg_{\mathcal{Q}} u_f(M') = u_f(M')$.

Now the fact that fully faithful functors reflect coproducts implies that $\text{im } u_f$ and w_f are an object respectively a morphism in \mathbf{Pos}_Π since $\tilde{\mathcal{O}}(\mathcal{P})$ has all coproducts and they are preserved by u_f . \square

Definition 4.20. Let $\check{\mathcal{S}} \in \mathbf{Pos}_\emptyset$, $\mathcal{S} \in \mathbf{Pos}_\Pi$, and $f \in \mathbf{Hom}_{\mathbf{Pos}_\emptyset}(\check{\mathcal{S}}, \mathcal{S})$. In this situation, we set

$$\mathcal{I}_s^f := w_f \downarrow s = \{i \in \text{im } u_f \mid w_f(i) \leq s\} \subseteq \text{im } u_f$$

which is a full subposet.

A reduced shape $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ is *boring* if there exists $s \in \mathcal{S}_{>\emptyset}$ such that $\mathcal{I}_s^\sigma = \{\emptyset\}$, i.e. such that for all $\check{s} \in \check{\mathcal{S}}$ with $\sigma(\check{s}) \leq s$ one has $\sigma(\check{s}) = \emptyset$.

Example 4.21. For any $n \in \mathbb{N}_0$, the shape \sqcup_n is reduced but not boring, since any $S \in \mathcal{P}(n)_{>\emptyset}$ has a subset of cardinality one.

Example 4.22. The inclusion of posets $\{0\} \rightarrow \{0 \leq 1\}$ is a boring reduced shape.

This definition is motivated by the statement of Proposition 4.24, which tells us that finite boring shapes do not have any interesting excision properties. The intuition is that, if there is an element $s \in \mathcal{S}$ with $\mathcal{I}_s^\sigma = \{\emptyset\}$, left Kan extension along σ copies the value at $0 \in \check{\mathcal{S}}$ to s (since s does not “see” anything else of $\check{\mathcal{S}}$), so that afterwards applying a functor F and taking the limit over $\mathcal{S}_{>\emptyset}$ yields F applied to that value again. However, before we can prove the formal statement, we need the following lemma.

Lemma 4.23. *Let $\check{\mathcal{S}} \in \mathbf{Pos}_\emptyset$, $\mathcal{S} \in \mathbf{Pos}_\Pi$, and $f \in \mathbf{Hom}_{\mathbf{Pos}_\emptyset}(\check{\mathcal{S}}, \mathcal{S})$. Then the poset \mathcal{I}_s^f has a terminal object.*

Proof. We claim that $i_* := \coprod_{i \in \mathcal{I}_s^f} i \in \mathbf{im} u_f$ is terminal in \mathcal{I}_s^f , where the coproduct is taken in $\mathbf{im} u_f$ and exists by Lemma 4.19. It is clear that we have $i \leq i_*$ for any $i \in \mathcal{I}_s^f$. But i_* also lies in \mathcal{I}_s^f as $w_f(i_*) = \coprod_{i \in \mathcal{I}_s^f} w_f(i) \leq s$ by Lemma 4.19 and the universal property of the coproduct. \square

Now we can prove that finite boring shapes are actually not interesting from an excision standpoint.

Proposition 4.24. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a finite boring reduced shape. Then any functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories is σ -excisive (as long as \mathcal{C} is σ -nice, and \mathcal{D} admits limits indexed by both $\mathcal{S}_{>\emptyset}$ and $(\mathbf{im} u_\sigma \times \mathcal{P})_{>\emptyset}$, where \mathcal{P} is as in the proof below).*

Proof. Let $D: \mathcal{S} \rightarrow \mathcal{C}$ be a σ -cocartesian diagram and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor. Furthermore, let \mathcal{P} be the full subposet $\{s \in \mathcal{S} \mid \mathcal{I}_s^\sigma = \{\emptyset\}\} \subseteq \mathcal{S}$. That σ is boring is then equivalent to \mathcal{P} having more elements than just \emptyset .

Now consider the diagram

$$\begin{array}{ccc}
 & (\mathbf{im} u_\sigma \times \mathcal{P})_{>\emptyset} & \\
 \text{pr} \swarrow & & \searrow c \\
 \mathbf{im} u_\sigma & \xrightarrow{\eta} & \mathcal{S}_{>\emptyset} \\
 \searrow w_\sigma & & \swarrow \text{inc} \\
 & \mathcal{S} &
 \end{array} \tag{4.2}$$

where inc denotes the inclusion, pr the (restriction of the) projection, c is given by $(i, p) \mapsto w_\sigma(i) \amalg p$, and the natural transformation η comes from the fact that $w_\sigma(i) \leq w_\sigma(i) \amalg p$.

Claim 4.24.1. *The functor c is homotopy initial.*

Proof. This is Lemma D.4 applied to w_σ and the inclusion $\text{inc}_\mathcal{P}: \mathcal{P} \rightarrow \mathcal{S}$. For this we need to show that for all $s \in \mathcal{S}_{>\emptyset}$ one of $w_\sigma \downarrow s$ and $\text{inc}_\mathcal{P} \downarrow s$ has a terminal object different from the respective initial object. Note that the category $w_\sigma \downarrow s = \mathcal{I}_s^\sigma$ always has a terminal object by Lemma 4.23. If this terminal object is equal to \emptyset then we have $s \in \mathcal{P}$ by definition, in which case $s \neq \emptyset$ is a terminal object of $\text{inc}_\mathcal{P} \downarrow s$. \blacksquare

Claim 4.24.2. *The functor pr is homotopy initial.*

Proof. We need to show that for all $i \in \mathbf{im} u_\sigma$ the category $\text{pr} \downarrow i$ is contractible. This category can be identified with $((\mathbf{im} u_\sigma) \downarrow i) \times \mathcal{P}_{>\emptyset}$. By Lemma D.3, it is now enough to show that \mathcal{P} has a terminal object different from \emptyset . For this we show that if $p, p' \in \mathcal{P}$ then also $p \amalg p' \in \mathcal{P}$ (where the coproduct is taken in \mathcal{S}), which is enough since \mathcal{P} has more than one element (by assumption) and is finite as it is a subset of \mathcal{S} . Otherwise

there would be $j \in (\text{im } u_\sigma)_{>\emptyset}$ such that $w_\sigma(j) \leq p \amalg p'$. Hence, by definition of u_σ , there is $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \neq \emptyset$ and $\sigma(\check{k}) \leq p \amalg p'$. Now we look at

$$\check{\mathcal{S}}_{p,p',\check{k}} = \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq p \text{ and } \sigma(\check{k}) \leq p' \amalg \sigma(\check{s})\}$$

which, since σ is a shape and $\sigma(\check{k}) \leq p \amalg p'$, is contractible. In particular it is non-empty, so let $\check{s} \in \check{\mathcal{S}}_{p,p',\check{k}}$. We have $\sigma(\check{s}) \leq p$, thus $\sigma(\check{s}) \in \mathcal{I}_p^\sigma$ (as $\sigma(\check{s}) \in \text{im } u_\sigma$), and hence $\sigma(\check{s}) = \emptyset$. Thus $\sigma(\check{k}) \leq p' \amalg \sigma(\check{s}) = p'$ and hence $\sigma(\check{k}) = \emptyset$, a contradiction. ■

Claim 4.24.3. *The natural transformation $D \circ \eta: D \circ w_\sigma \circ \text{pr} \rightarrow D \circ \text{inc} \circ c$ is an equivalence.*

Proof. Since D is σ -cocartesian, we have $D \simeq \text{Lan}_\sigma \text{Res}_\sigma D$. By Lemma C.5 the transformation $(\text{Lan}_\sigma \text{Res}_\sigma)(D) \circ \eta$ is given, at $(i, p) \in (\text{im } u_\sigma \times \mathcal{P})_{>\emptyset}$, by the map

$$\text{colim}_{\sigma \downarrow w_\sigma(i)} \left(\text{Res}_\sigma(D) \circ \text{pr}_{\sigma \downarrow w_\sigma(i)} \right) \longrightarrow \text{colim}_{\sigma \downarrow (w_\sigma(i) \amalg p)} \left(\text{Res}_\sigma(D) \circ \text{pr}_{\sigma \downarrow (w_\sigma(i) \amalg p)} \right)$$

induced by the canonical functor $\sigma \downarrow w_\sigma(i) \rightarrow \sigma \downarrow (w_\sigma(i) \amalg p)$.

We will now show that $\sigma \downarrow w_\sigma(i) \rightarrow \sigma \downarrow (w_\sigma(i) \amalg p)$ is an isomorphism, for which surjectivity (on objects) suffices as the map is an inclusion of full subposets of $\check{\mathcal{S}}$. For this we need that, if $\check{k} \in \check{\mathcal{S}}$ such that $\sigma(\check{k}) \leq w_\sigma(i) \amalg p$, then already $\sigma(\check{k}) \leq w_\sigma(i)$. For this consider

$$\check{\mathcal{S}}_{p,w_\sigma(i),\check{k}} = \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq p \text{ and } \sigma(\check{k}) \leq w_\sigma(i) \amalg \sigma(\check{s})\}$$

which is contractible (since $\sigma(\check{k}) \leq w_\sigma(i) \amalg p$), hence non-empty. So let \check{s} be an element of $\check{\mathcal{S}}_{p,w_\sigma(i),\check{k}}$. The condition $\sigma(\check{s}) \leq p$ implies $\sigma(\check{s}) = \emptyset$, thus we have $\sigma(\check{k}) \leq w_\sigma(i) \amalg \emptyset = w_\sigma(i)$, which we wanted to show. ■

Now we note that diagram (4.2) extends to a diagram

$$\begin{array}{ccc} & ((\text{im } u_\sigma \times \mathcal{P})_{>\emptyset})^\triangleleft & \\ \text{pr}^\triangleleft \swarrow & & \searrow c^\triangleleft \\ (\text{im } u_\sigma)^\triangleleft & \xrightarrow{\eta^\triangleleft} & (\mathcal{S}_{>\emptyset})^\triangleleft \\ (w_\sigma)^\triangleleft \swarrow & & \searrow \text{inc}^\triangleleft \\ & \mathcal{S}^\triangleleft & \\ & \downarrow q & \\ & \mathcal{S} & \end{array}$$

where η^\triangleleft is given by the identity at the cone point and by η otherwise, and q is the identity on \mathcal{S} and \emptyset at the cone point. This allows us to obtain maps from $F(D(\emptyset))$

into certain limits indexed by the categories occurring in diagram (4.2). Namely we get, by (the dual of) Lemma C.9, the following homotopy commutative diagram

$$\begin{array}{ccc}
\lim_{(\text{im } u_\sigma \times \mathcal{P})_{> \emptyset}} (F \circ D \circ w_\sigma \circ \text{pr}) & \xrightarrow{\simeq_\eta} & \lim_{(\text{im } u_\sigma \times \mathcal{P})_{> \emptyset}} (F \circ D \circ \text{inc} \circ c) \\
\uparrow \text{pr}^* \simeq & \swarrow & \nearrow \\
& F(D(\emptyset)) & \\
\uparrow \simeq & \nwarrow & \searrow \\
\lim_{\text{im } u_\sigma} (F \circ D \circ w_\sigma) & & \lim_{\mathcal{S}_{> \emptyset}} (F \circ D \circ \text{inc}) \\
& & \uparrow \simeq c^*
\end{array}$$

where the maps around the boundary are equivalences by the previous three claims and the map into the bottom left is one by Lemma C.10 as \emptyset is initial in $\text{im } u_\sigma$ and $q \circ (w_\sigma)^\triangleleft$ applied to the unique map from the cone point to \emptyset is the identity. Hence the map into the bottom right is an equivalence as well, which is what we wanted to show. \square

The rest of this subsection is devoted to comparing a (pre)shape $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ to the free shape $o_{\check{\mathcal{S}}}: \check{\mathcal{S}} \rightarrow \tilde{\mathcal{O}}(\check{\mathcal{S}})$.

Proposition 4.25. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full preshape. Then we have a map of posets $e: \mathcal{S} \rightarrow \tilde{\mathcal{O}}(\check{\mathcal{S}})$ defined by $s \mapsto \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s\}$. If $e^{-1}(\{\emptyset_{\mathcal{S}}\}) = \{\emptyset_{\check{\mathcal{S}}}\}$, this gives a map of preshapes*

$$\begin{array}{ccc}
\check{\mathcal{S}} & \xrightarrow{\text{id}} & \check{\mathcal{S}} \\
\sigma \downarrow & & \downarrow o_{\check{\mathcal{S}}} \\
\mathcal{S} & \xrightarrow{e} & \tilde{\mathcal{O}}(\check{\mathcal{S}})
\end{array}$$

that is indirect. Furthermore, if we additionally assume σ to be a shape, the condition $e^{-1}(\{\emptyset_{\mathcal{S}}\}) = \{\emptyset_{\check{\mathcal{S}}}\}$ is equivalent to σ being non-boring.

Proof. For commutativity of the square we need, for all $\check{k} \in \check{\mathcal{S}}$, the equality $\{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq \sigma(\check{k})\} = \{\check{s} \in \check{\mathcal{S}} \mid \check{s} \leq \check{k}\}$, which follows from σ being full. For indirectness, note that, for all $s \in \mathcal{S}$, the induced functor $\sigma \downarrow s \rightarrow o_{\check{\mathcal{S}}} \downarrow e(s)$ is an isomorphism since both sides can be identified with the full subposet $e(s) = \{\check{s} \mid \sigma(\check{s}) \leq s\} \subseteq \check{\mathcal{S}}$. For the last statement of the proposition, note that σ being non-boring is equivalent to, for all $s \in \mathcal{S}_{> \emptyset}$, there existing an $\check{s} \in \check{\mathcal{S}}$ such that $\emptyset_{\mathcal{S}} \neq \sigma(\check{s}) \leq s$, which is a reformulation of the statement $e^{-1}(\{\emptyset_{\mathcal{S}}\}) = \{\emptyset_{\check{\mathcal{S}}}\}$. \square

Corollary 4.26. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a non-boring full shape. Then a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories that is σ -excisive is also $o_{\check{\mathcal{S}}}$ -excisive (as long as \mathcal{C} is σ -nice and $o_{\check{\mathcal{S}}}$ -nice, and \mathcal{D} admits limits indexed both by $\mathcal{S}_{> \emptyset}$ and by $\tilde{\mathcal{O}}(\check{\mathcal{S}})_{> \emptyset}$).*

Combining this with Corollary 4.17, we obtain:

Theorem 4.27. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a finite non-boring full shape. Then a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories that is σ -excisive is also $(n-1)$ -excisive, where n is the number of minimal elements of $\check{\mathcal{S}}_{>\emptyset}$ (as long as \mathcal{C} admits all finite colimits, and \mathcal{D} admits limits indexed by $\mathcal{P}(n)_{>\emptyset}$, $\mathcal{S}_{>\emptyset}$, and $\tilde{\mathcal{O}}(\check{\mathcal{S}})_{>\emptyset}$).*

The next two results tell us that any non-boring full shape $\check{\mathcal{S}} \rightarrow \mathcal{S}$ has the same excision properties as a shape $\check{\mathcal{S}} \rightarrow \mathcal{Q}$ such that \mathcal{Q} is a retract of $\tilde{\mathcal{O}}(\check{\mathcal{S}})$. In particular, for a fixed finite $\check{\mathcal{S}}$, there are only finitely many full shapes of the form $\check{\mathcal{S}} \rightarrow \mathcal{S}$ with different excision properties. Combining this with Proposition 4.7, we even get this result if we do not require the shapes to be full. See Corollary 5.17 for a precise version of this statement.

Proposition 4.28. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced shape. Then $v_\sigma: \check{\mathcal{S}} \rightarrow \text{im } u_\sigma$ (see Definition 4.18) is again a shape which is finite if σ is, and the map of shapes*

$$\begin{array}{ccc} \check{\mathcal{S}} & \xrightarrow{\text{id}} & \check{\mathcal{S}} \\ v_\sigma \downarrow & & \downarrow \sigma \\ \text{im } u_\sigma & \xrightarrow{w_\sigma} & \mathcal{S} \end{array}$$

is indirect. Furthermore, if σ is full and non-boring, then it is also direct.

Proof. The first part follows from Lemma 3.7 using Lemma 4.19. For the second part we will use the following claim:

Claim 4.28.1. *Let the functor $r: \mathcal{S} \rightarrow \text{im } u_\sigma$ be given by $s \mapsto \max \mathcal{I}_s^\sigma$ and assume it restricts to a functor $\mathcal{S}_{>\emptyset} \rightarrow (\text{im } u_\sigma)_{>\emptyset}$. Then $r|_{\mathcal{S}_{>\emptyset}}$ is right adjoint to $w_\sigma|_{(\text{im } u_\sigma)_{>\emptyset}}$.*

Proof. First note that $\max \mathcal{I}_s^\sigma$ exists by Lemma 4.23 and that r is functorial as $s \leq s'$ implies $\mathcal{I}_s^\sigma \subseteq \mathcal{I}_{s'}^\sigma$ and hence $\max \mathcal{I}_s^\sigma \leq \max \mathcal{I}_{s'}^\sigma$. For the adjunction we have to show that, for all $i \in (\text{im } u_\sigma)_{>\emptyset}$ and $s \in \mathcal{S}_{>\emptyset}$, we have $i \leq r(s)$ if and only if $w_\sigma(i) \leq s$, which follows from \mathcal{I}_s^σ being defined as $w_\sigma \downarrow s$. \blacksquare

Note that our assumption that σ is not boring precisely says that r restricts to a functor $\mathcal{S}_{>\emptyset} \rightarrow (\text{im } u_\sigma)_{>\emptyset}$. Hence $w_\sigma|_{(\text{im } u_\sigma)_{>\emptyset}}: (\text{im } u_\sigma)_{>\emptyset} \rightarrow \mathcal{S}_{>\emptyset}$ is left adjoint and thus homotopy initial. Together with w_σ and σ being full this shows directness \square

Proposition 4.29. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full shape and assume that $u_\sigma: \tilde{\mathcal{O}}(\check{\mathcal{S}}) \rightarrow \mathcal{S}$ is surjective. Then $u_\sigma \circ e = \text{id}_\mathcal{S}$, where e is as in Proposition 4.25. In particular σ is a retract of $o_{\check{\mathcal{S}}}$ (in the category $\check{\mathcal{S}} \downarrow \mathbf{Pos}_\emptyset$).*

Proof. We need to show that $u_\sigma(e(s)) = s$ for all $s \in \mathcal{S}$, where $e(s) = \{\check{s} \in \check{\mathcal{S}} \mid \sigma(\check{s}) \leq s\}$. So, let s be an element of \mathcal{S} . By assumption, there is an $M \subseteq \check{\mathcal{S}}$ such that $s = u_\sigma(M) = \coprod_{m \in M} \sigma(m)$. This implies $\sigma(m) \leq s$ for all $m \in M$, hence $M \subseteq e(s)$. We obtain the inequalities $s = u_\sigma(M) \leq u_\sigma(e(s)) = \coprod_{\check{s} \in e(s)} \sigma(\check{s}) \leq s$, which imply $u_\sigma(e(s)) = s$. \square

Lemma 4.30. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a reduced shape. Then the following diagram commutes*

$$\begin{array}{ccc}
 \tilde{\mathcal{O}}(\check{\mathcal{S}}) & \xrightarrow{u_\sigma} & \mathcal{S} \\
 & \searrow^{u_{v_\sigma}} & \nearrow^{w_f} \\
 & \text{im } u_\sigma &
 \end{array}$$

which in particular implies that u_{v_σ} is surjective (as u_σ maps surjectively onto $\text{im } u_\sigma$).

Proof. This follows directly from Lemma 4.19. □

Theorem 4.31. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a full shape. Then $v_\sigma: \check{\mathcal{S}} \rightarrow \text{im } u_\sigma$ is a retract of $o_{\check{\mathcal{S}}}$ (in the category $\check{\mathcal{S}} \downarrow \mathbf{Pos}_\emptyset$). Moreover, if σ is non-boring, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between ∞ -categories is σ -excisive if and only if it is v_σ -excisive (as long as \mathcal{C} is σ -nice (which implies v_σ -nice), and \mathcal{D} admits limits indexed by both $\mathcal{S}_{>\emptyset}$ and $(\text{im } u_\sigma)_{>\emptyset}$).*

Proof. This follows by applying Proposition 4.28 to σ and then Proposition 4.29 to v_σ . For the latter part we use that v_σ is full by Lemma D.2 since σ is, and that u_{v_σ} is actually surjective by Lemma 4.30. □

5. The Taylor graph

The goal of this section is to assemble all of the various excisive approximations for different (pre)shapes into a natural coherent diagram lying under the functor we are approximating, analogous to the Taylor tower in classical Goodwillie calculus. For this we first have to define the category this will be indexed by. Since the existences of the approximations depend on the ∞ -categories \mathcal{C} and \mathcal{D} between which we consider functors, the indexing category also needs to depend on these ∞ -categories. However, later in this section, we will also obtain a version which puts more conditions on \mathcal{C} and \mathcal{D} but in return uses a fixed indexing category.

Definition 5.1. Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape and \mathcal{C} and \mathcal{D} two ∞ -categories. We say that $\text{Fun}(\mathcal{C}, \mathcal{D})$ admits a universal σ -excisive approximation if the inclusion $\text{inc}: \text{Exc}_\sigma(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ has a left adjoint. In this case we fix an adjunction $P_\sigma \dashv \text{inc}$ and denote its unit by $p_\sigma: \text{id} \rightarrow \text{inc} \circ P_\sigma$.

Remark 5.2. By Theorem 2.42, we can, for σ a full shape, choose P_σ and p_σ so that they agree with our previous definition (at least when \mathcal{C} and \mathcal{D} are nice enough). In the other cases this replaces the old notion, as it will not be useful to us anymore.

Notation 5.3. Let \mathcal{C} and \mathcal{D} be ∞ -categories. We write $\mathbf{PreSha}_{\mathcal{C}, \mathcal{D}}$ for the thin category with objects the preshapes σ such that $\text{Fun}(\mathcal{C}, \mathcal{D})$ admits a universal σ -excisive approximation and such that there is a morphism from σ to τ if and only if any functor $F: \mathcal{C} \rightarrow \mathcal{D}$ that is σ -excisive is also τ -excisive.

The main input in the construction of these diagrams is the following lemma. Both statement and proof are due to Markus Land (though any mistake is mine).

Lemma 5.4. Let \mathcal{I} be a category and \mathcal{C} an ∞ -category. Denote by $\mathcal{I}_{\text{disc}} \subseteq \mathcal{I}$ the discrete subcategory containing all objects and let $f: \mathcal{I}_{\text{disc}} \rightarrow \mathcal{C}$ be a functor such that, for all x and y in \mathcal{I} such that there is a morphism $x \rightarrow y$, the mapping space $\text{Map}_{\mathcal{C}}(f(x), f(y))$ is contractible. Then there is an essentially unique extension of f to a functor $\mathcal{I} \rightarrow \mathcal{C}$.

Proof. We want to show that the pullback \mathcal{D} (in the 1-category of simplicial sets) of the lower right corner of the middle part of the diagram

$$\begin{array}{ccccc}
 \partial\Delta^n & \xrightarrow{s} & \mathcal{D} & \xrightarrow{p} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
 \downarrow & \nearrow \text{dashed} & \downarrow & & \downarrow \text{Res}_{\text{inc}} \\
 \Delta^n & \longrightarrow & * & \xrightarrow{\text{const}_f} & \text{Fun}(\mathcal{I}_{\text{disc}}, \mathcal{C})
 \end{array}$$

is a contractible Kan complex, i.e. that any map $s: \partial\Delta^n \rightarrow \mathcal{D}$ extends to a map $\Delta^n \rightarrow \mathcal{D}$ as indicated on the left side of the above diagram. By the universal property of the pullback, this is equivalent to finding a map $t: \Delta^n \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$ such that $t \circ \text{inc}_{\partial\Delta^n} = p \circ s$ and $\text{Res}_{\text{inc}} \circ t = \text{const}_f$. By currying t corresponds to a map $\Delta^n \times \mathcal{I} \rightarrow \mathcal{C}$ such that it

restricts to $f \circ \text{pr}_2$ on $\Delta^n \times \mathcal{I}_{\text{disc}}$ and to the curried morphism associated to $p \circ s$ on $\partial\Delta^n \times \mathcal{I}$. Said differently, we want to find a dashed extension as in the following diagram

$$\begin{array}{ccc} \Delta^n \times \mathcal{I}_{\text{disc}} \cup \partial\Delta^n \times \mathcal{I} & \xrightarrow{l} & \mathcal{C} \\ \downarrow & \dashrightarrow & \\ \Delta^n \times \mathcal{I} & & \end{array}$$

where l is given by $f \circ \text{pr}_2$ on $\Delta^n \times \mathcal{I}_{\text{disc}}$ and by $p' \circ (s \times \text{id}_{\mathcal{I}})$ on $\partial\Delta^n \times \mathcal{I}$, where $p': \mathcal{D} \times \mathcal{I} \rightarrow \mathcal{C}$ is the curried morphism associated to p . We will now factor l through another ∞ -category that only sees the information relevant to us and we can thus control.

For this, note that our assumptions imply the existence of a unique extension g of the composition $\pi_{\mathcal{C}} \circ f$ to \mathcal{I} (here $\pi_{\mathcal{C}}$ is the canonical map $\mathcal{C} \rightarrow \text{h}\mathcal{C}$). Now we write \mathcal{E} for the pullback (in the 1-category of simplicial sets) of the lower right corner in the following diagram

$$\begin{array}{ccccc} \mathcal{I}_{\text{disc}} & & & & \\ & \xrightarrow{f} & & & \\ & \hat{f} \dashrightarrow & \mathcal{E} & \xrightarrow{k} & \mathcal{C} \\ & \text{inc} \searrow & \downarrow q & & \downarrow \pi_{\mathcal{C}} \\ & & \mathcal{I} & \xrightarrow{g} & \text{h}\mathcal{C} \end{array}$$

and note that, since the large outer diagram commutes, we obtain a map \hat{f} making the two triangles commute (in particular f factors through \mathcal{E}). Now, by Lemma D.7, we obtain that the pullback \mathcal{E} is again an ∞ -category and that q can be identified with $\pi_{\mathcal{E}}$, so in particular that q is essentially surjective. Using our condition on f , the lemma also implies that q is fully faithful, hence a categorical equivalence. (Note that we can not just use the subcategory spanned by the essential image $\text{im } g$ instead of \mathcal{E} since, when g is not full, there could be morphisms in $\text{im } g$ that our conditions do not control.)

To see that l actually factors through \mathcal{E} , we will first show that p' factors through \mathcal{E} . For this, consider the commutative diagram

$$\begin{array}{ccccc} \mathcal{D} & \xrightarrow{p} & \text{Fun}(\mathcal{I}, \mathcal{C}) & \xrightarrow{\pi_{\mathcal{C}} \circ} & \text{Fun}(\mathcal{I}, \text{h}\mathcal{C}) \\ \downarrow & & \text{Res}_{\text{inc}} \downarrow & & \downarrow \text{Res}_{\text{inc}} \\ * & \xrightarrow{\text{const}_f} & \text{Fun}(\mathcal{I}_{\text{disc}}, \mathcal{C}) & \xrightarrow{\pi_{\mathcal{C}} \circ} & \text{Fun}(\mathcal{I}_{\text{disc}}, \text{h}\mathcal{C}) \end{array}$$

and note that $(\pi_{\mathcal{C}} \circ) \circ p$ is just const_g since the subcategory of $\text{Fun}(\mathcal{I}, \text{h}\mathcal{C})$ lying over (the discrete subcategory spanned by) the object $\pi_{\mathcal{C}} \circ f \in \text{Fun}(\mathcal{I}_{\text{disc}}, \text{h}\mathcal{C})$ has the single object

g and is discrete. Hence p' fits into a commutative diagram of the form

$$\begin{array}{ccc}
\mathcal{D} \times \mathcal{I}_{\text{disc}} & & \mathcal{C} \\
\downarrow \text{id}_{\mathcal{D}} \times \text{inc} & \nearrow f \circ \text{pr}_2 & \downarrow \pi_{\mathcal{C}} \\
\mathcal{D} \times \mathcal{I} & \xrightarrow{p'} & \mathcal{C} \\
\downarrow \text{pr}_{\mathcal{I}} & & \downarrow \pi_{\mathcal{C}} \\
\mathcal{I} & \xrightarrow{g} & \text{h}\mathcal{C}
\end{array}$$

$\text{inc} \circ \text{pr}_2$ (curved arrow from $\mathcal{D} \times \mathcal{I}_{\text{disc}}$ to \mathcal{I})

and we obtain a functor $\hat{p}': \mathcal{D} \times \mathcal{I} \rightarrow \mathcal{E}$ such that $k \circ \hat{p}' = p'$ and $q \circ \hat{p}' = \text{pr}_{\mathcal{I}}$. Thus also $k \circ \hat{p}' \circ (\text{id}_{\mathcal{D}} \times \text{inc}) = f \circ \text{pr}_2$ and $q \circ \hat{p}' \circ (\text{id}_{\mathcal{D}} \times \text{inc}) = \text{inc} \circ \text{pr}_2$, which implies $\hat{p}' \circ (\text{id}_{\mathcal{D}} \times \text{inc}) = \hat{f} \circ \text{pr}_2$ by the universal property of \mathcal{E} .

Now we can construct the following diagram

$$\begin{array}{ccc}
\Delta^n \times \mathcal{I}_{\text{disc}} \cup \partial\Delta^n \times \mathcal{I} & \xrightarrow{\hat{l}} & \mathcal{E} & \xrightarrow{k} & \mathcal{C} \\
\downarrow & \nearrow \text{dashed} & \downarrow q & & \\
\Delta^n \times \mathcal{I} & \xrightarrow{\text{pr}_{\mathcal{I}}} & \mathcal{I} & &
\end{array}$$

where \hat{l} is given by $\hat{f} \circ \text{pr}_2$ on $\Delta^n \times \mathcal{I}_{\text{disc}}$ and by $\hat{p}' \circ (s \times \text{id}_{\mathcal{I}})$ on $\partial\Delta^n \times \mathcal{I}$. It follows from what we said before that \hat{l} is well-defined, that the diagram commutes (without the dashed arrow), and that we actually have $k \circ \hat{l} = l$. Remembering that q is a categorical equivalence and, by Lemma D.8, also a categorical fibration, we obtain the desired dashed lift since trivial categorical fibrations have the right lifting property against inclusions of simplicial sets. \square

To use the above lemma we need information about mapping spaces in slice categories (since we want a diagram which lies under a given functor in a coherent way) which the following lemma and its consequences will provide.

Lemma 5.5. *Let \mathcal{C} be an ∞ -category and $i: \mathcal{C}_0 \rightarrow \mathcal{C}$ the inclusion of a full subcategory such that there is an adjunction $l \dashv i$ with unit $\eta: \text{id} \rightarrow i \circ l$. Further let $C \in \mathcal{C}$, $C_0 \in \mathcal{C}_0$, and $f: C \rightarrow i(C_0)$ a morphism in \mathcal{C} . Then $\text{Map}_{\mathcal{C}_{C_0}}(\eta(C), f)$ is contractible.*

Proof. By [LurHTT, Lemma 5.5.5.12], we have that $\text{Map}_{\mathcal{C}_{C_0}}(\eta(C), f)$ is a homotopy fiber of the map $(\circ \eta(C)): \text{Map}_{\mathcal{C}}(i(l(C)), i(C_0)) \rightarrow \text{Map}_{\mathcal{C}}(C, i(C_0))$ over the point f . But, by Lemma D.10, the composition

$$\text{Map}_{\mathcal{C}_0}(l(C), C_0) \xrightarrow{i} \text{Map}_{\mathcal{C}}(i(l(C)), i(C_0)) \xrightarrow{\circ \eta(C)} \text{Map}_{\mathcal{C}}(C, i(C_0))$$

is an equivalence. Now we note that the first map in this composition is an equivalence since i is fully faithful. Hence the second map is also an equivalence and thus the homotopy fiber we are interested in is trivial. \square

Corollary 5.6. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape and \mathcal{C} and \mathcal{D} ∞ -categories such that $\text{Fun}(\mathcal{C}, \mathcal{D})$ admits a universal σ -excisive approximation. Furthermore, let $\alpha: F \rightarrow G$ be a natural transformation of functors $\mathcal{C} \rightarrow \mathcal{D}$ such that G is σ -excisive. Then $\text{Map}_{\text{Fun}(\mathcal{C}, \mathcal{D})_{F/}}(\text{p}_\sigma(F), \alpha)$ is contractible.*

Lemma 5.7. *Let \mathcal{C} be an ∞ -category and $i: \mathcal{C}_0 \rightarrow \mathcal{C}$ the inclusion of a full subcategory such that there is an adjunction $l \dashv i$ with unit $\eta: \text{id} \rightarrow i \circ l$. Furthermore, let \mathcal{E} be an ∞ -category, $F: \mathcal{E} \rightarrow \mathcal{C}$ and $F_0: \mathcal{E} \rightarrow \mathcal{C}_0$ functors, and $\alpha: F \rightarrow i \circ F_0$ a natural transformation. Then $\text{Map}_{\text{Fun}(\mathcal{E}, \mathcal{C})_{F/}}(\eta \circ F, \alpha)$ is contractible.*

Proof. By Lemma D.5, there is an adjunction with left adjoint $(l \circ): \text{Fun}(\mathcal{E}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{E}, \mathcal{C}_0)$, right adjoint $(i \circ): \text{Fun}(\mathcal{E}, \mathcal{C}_0) \rightarrow \text{Fun}(\mathcal{E}, \mathcal{C})$, and unit $(\eta \circ)$. Noting that $(i \circ)$ is the inclusion of a full subcategory, we can apply Lemma 5.5 to obtain the desired statement. \square

Corollary 5.8. *Let $\sigma: \check{\mathcal{S}} \rightarrow \mathcal{S}$ be a preshape and \mathcal{C} and \mathcal{D} ∞ -categories such that $\text{Fun}(\mathcal{C}, \mathcal{D})$ admits a universal σ -excisive approximation. Furthermore, let $A: \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Exc}_\sigma(\mathcal{C}, \mathcal{D})$ be a functor and $\alpha: \text{id}_{\text{Fun}(\mathcal{C}, \mathcal{D})} \rightarrow \text{inc} \circ A$ a natural transformation. Then the space of maps from p_σ to α in $\text{Fun}(\text{Fun}(\mathcal{C}, \mathcal{D}), \text{Fun}(\mathcal{C}, \mathcal{D}))_{\text{id}/}$ is contractible.*

We now know enough to be able to construct a diagram as promised in the beginning of this section. There are two versions: one which also takes into account maps between functors (i.e. makes the naturality of the diagram precise), and a second one only considering a single functor.

Theorem 5.9. *Let \mathcal{C} and \mathcal{D} be two ∞ -categories.*

a) *There is an essentially unique functor*

$$\text{P}: (\mathbf{PreSha}_{\mathcal{C}, \mathcal{D}})^{\text{op}} \longrightarrow \text{Fun}(\text{Fun}(\mathcal{C}, \mathcal{D}), \text{Fun}(\mathcal{C}, \mathcal{D}))_{\text{id}/}$$

such that $\text{P}(\sigma) = \text{p}_\sigma$.

b) *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor. Then there is an essentially unique functor*

$$\text{P}(F): (\mathbf{PreSha}_{\mathcal{C}, \mathcal{D}})^{\text{op}} \longrightarrow \text{Fun}(\mathcal{C}, \mathcal{D})_{F/}$$

such that $\text{P}(F)(\sigma) = \text{p}_\sigma(F)$.

In particular, we obtain such functors for any subcategory of $\mathbf{PreSha}_{\mathcal{C}, \mathcal{D}}$. These restricted functors are also essentially unique.

Proof. We want to apply Lemma 5.4 to obtain the desired functors. For the first point, note that, if there is a map $\sigma \rightarrow \tau$ in $\mathbf{PreSha}_{\mathcal{C}, \mathcal{D}}$, then P_σ lands in $\text{Exc}_\tau(\mathcal{C}, \mathcal{D})$, hence Corollary 5.8 implies that $\text{Map}(\text{p}_\tau, \text{p}_\sigma)$ is contractible. The second follows in the same way, using Corollary 5.6 instead of Corollary 5.8. The essential uniqueness of the restrictions also follows from Lemma 5.4. \square

Remark 5.10. Using the essential uniqueness of $P(F)$, we see that it is equivalent to the functor obtained from P by postcomposing with evaluation at F , which explains our notation.

We now describe a version of these diagrams with a single indexing category independent of the ∞ -categories \mathcal{C} and \mathcal{D} .

Notation 5.11. We write $\mathbf{Sha}_{\text{fin}}$ for the thin (i.e. each hom-set has at most one element) category with

- objects the collection of (small) finite shapes
- a morphism from σ to τ if and only if, for all ∞ -categories \mathcal{C} and \mathcal{D} such that \mathcal{C} has a terminal object and admits all finite colimits and such that \mathcal{D} is differentiable, each functor $F: \mathcal{C} \rightarrow \mathcal{D}$ that is σ -excisive is also τ -excisive.

Definition 5.12. We will say that two finite shapes are *equivalent* if they are isomorphic in $\mathbf{Sha}_{\text{fin}}$.

Corollary 5.13. *Let \mathcal{C} be an ∞ -category that has a terminal object and admits all finite colimits, and \mathcal{D} a differentiable ∞ -category.*

a) *There is an essentially unique functor*

$$P: (\mathbf{Sha}_{\text{fin}})^{\text{op}} \longrightarrow \text{Fun}(\text{Fun}(\mathcal{C}, \mathcal{D}), \text{Fun}(\mathcal{C}, \mathcal{D}))_{\text{id}/}$$

such that $P(\sigma) = p_\sigma$.

b) *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor. Then there is an essentially unique functor*

$$P(F): (\mathbf{Sha}_{\text{fin}})^{\text{op}} \longrightarrow \text{Fun}(\mathcal{C}, \mathcal{D})_{F/}$$

such that $P(F)(\sigma) = p_\sigma(F)$.

In particular, we obtain such functors for any subcategory of $\mathbf{Sha}_{\text{fin}}$. These restricted functors are also essentially unique.

Proof. By Corollary 4.8, there is a functorial inclusion $\mathbf{Sha}_{\text{fin}} \rightarrow \mathbf{PreSha}_{\mathcal{C}, \mathcal{D}}$. Then Theorem 5.9 implies the statement. \square

Remark 5.14. There are analogous statements for shapes with higher cardinality bounds (as long as we restrict ourselves to full shapes). However, they are less useful as the necessary differentiability condition on \mathcal{D} becomes very strong.

5.1. Relation to the Taylor tower

Notation 5.15. We write $\mathbf{Sha}_{\text{fin,nb}}$ for the full (thin) subcategory of $\mathbf{Sha}_{\text{fin}}$ spanned by the shapes equivalent to a finite non-boring full shape.

The following proposition tells us that restricting ourselves to $\mathbf{Sha}_{\text{fin,nb}}$ drops precisely the shapes with uninteresting excision properties.

Proposition 5.16. *A finite shape σ is equivalent to a finite boring full shape if and only if, for all ∞ -categories \mathcal{C} and \mathcal{D} such that \mathcal{C} has a terminal object and admits all finite colimits and such that \mathcal{D} is differentiable, each functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is σ -excisive. Furthermore, a finite non-boring full shape is not equivalent to a finite boring full shape.*

In particular, the finite shapes equivalent to a finite boring full shape are precisely the terminal objects of $\mathbf{Sha}_{\text{fin}}$, and $\mathbf{Sha}_{\text{fin,nb}}$ consists precisely of the non-terminal objects of $\mathbf{Sha}_{\text{fin}}$.

Proof. By Proposition 4.24, any shape equivalent to a finite boring full shape has trivial excision properties. This shows one direction of the first statement.

Moreover, by Theorem 4.27, for any $\sigma \in \mathbf{Sha}_{\text{fin,nb}}$, there exists an $n \in \mathbb{N}_0$ such that σ -excisive implies n -excisive. Since being n -excisive is a non-trivial condition (i.e. there exists a functor $\mathcal{C} \rightarrow \mathcal{D}$ (with \mathcal{C} and \mathcal{D} as above) that is not n -excisive; one example is the functor F from pointed spaces to spectra given by $X \mapsto \Sigma^\infty(X^{\wedge(n+1)})$ as it is not weakly constant but still $(n+1)$ -reduced, i.e. $P_n F$ is terminal, see [Goo03, Remark 1.16]), we obtain that being σ -excisive is also a non-trivial condition. In particular, any shape in $\mathbf{Sha}_{\text{fin,nb}}$ can not be equivalent to a finite boring full shape. This shows the second statement.

For the other implication of the first statement, note that any finite shape is equivalent to a finite full shape by Proposition 4.7. But, if it has trivial excision properties, it can't be equivalent to a finite non-boring full shape by the same argument as for the second statement. So it must be equivalent to a finite boring full shape.

The last statement now follows by noting that, since there exists a finite shape σ such that any functor is σ -excisive (see Example 4.22), these shapes are precisely the terminal objects of $\mathbf{Sha}_{\text{fin}}$. \square

We also have the following direct corollary of Theorem 4.31 and Proposition 4.7, which tells us a bit more about the structure of $\mathbf{Sha}_{\text{fin,nb}}$:

Corollary 5.17. *Let $\tilde{\mathcal{S}}$ be a finite poset. Then the full subcategory of $\mathbf{Sha}_{\text{fin,nb}}$ spanned by the shapes with domain $\tilde{\mathcal{S}}$ (i.e. those of the form $\sigma: \tilde{\mathcal{S}} \rightarrow \mathcal{S}$ for some \mathcal{S}) has only finitely many isomorphism classes.*

We will now state and prove the second main theorem of this thesis, relating our class of finite non-boring shapes to the original cubes.

Theorem 5.18. *The functor $C: (\mathbb{N}_0)^{\text{op}} \rightarrow (\mathbf{Sha}_{\text{fin,nb}})^{\text{op}}$ given by sending n to \llcorner_{n+1} is homotopy initial.*

Proof. First note that C is well-defined since \sqcup_{n+1} is a shape (see Example 2.37) that is full but not boring (see Example 4.21) and since n -excisive implies $(n+1)$ -excisive (see Corollary 3.6). To see that it is homotopy initial we need to show that, for all $\sigma \in (\mathbf{Sha}_{\text{fin,nb}})^{\text{op}}$, the category $C \downarrow \sigma$ is contractible. However, this is just the full subposet $\mathcal{P} \subseteq (\mathbb{N}_0)^{\text{op}}$ spanned by those n such that σ -excisive implies n -excisive. By definition of $\mathbf{Sha}_{\text{fin,nb}}$ we can assume that σ is full and non-boring. Hence, by Theorem 4.27, the poset \mathcal{P} is not empty, and, by Corollary 3.6, it is closed below (i.e. if it contains n then it contains all m such that $m \leq n$ in $(\mathbb{N}_0)^{\text{op}}$). But any non-empty subposet of $(\mathbb{N}_0)^{\text{op}}$ that is downward closed is isomorphic to $(\mathbb{N}_0)^{\text{op}}$, hence contractible (since it has a terminal object). \square

Remark 5.19. This theorem tells us in particular that the Taylor graph $P(F)$ of a functor F converges if and only if its Taylor tower $P(F) \circ C$ does (here convergence means that the canonical map from F to the limit of the respective diagram is an equivalence). In particular any convergence criteria for the tower, such as the analytic functors of Goodwillie (cf. [Goo92, Definition 4.2] and [Goo03, Theorem 1.13]), can also be applied to the graph.

Theorem 5.18 suggests the following question, which is, to the authors knowledge, still unresolved:

Open Question 5.20. *Is the functor C an equivalence of categories? Or, phrased differently, is any shape in $\mathbf{Sha}_{\text{fin,nb}}$ equivalent to \sqcup_n for some $n \in \mathbb{N}_0$?*

Should this be the case, the Taylor graph P would not contain more information than the classical Taylor tower $P \circ C$, which would be evidence that the cubes are the correct shapes to consider. Otherwise the Taylor graph could give a potentially interesting, finer resolution of the tower.

A. The calculus of mates

In this appendix we recall the mate construction as well as a number of lemmas concerning it, which are quite useful when working with adjunctions and natural transformations. Since this is not supposed to be a comprehensive exposition of the topic, we will be brief and only state and give references for the statements we will use. A concise summary of these, and a few more, important statements, though without proofs, can be found in [GPS, Appendix A]. A longer exposition with proofs is given (in French) in [Ayo, Section 1.1.2].

Notation A.1. Suppose we have, in a (strict) 2-category, a diagram of the form

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ h \downarrow & \alpha \swarrow & \downarrow k \\ C & \xrightarrow{c} & D \end{array}$$

and fixed adjunctions $a_! \dashv a$ and $c_! \dashv c$. In this situation we write $\alpha_!$ for the *mate* of α , which is a 2-morphism of the form

$$\begin{array}{ccc} A & \xleftarrow{a_!} & B \\ h \downarrow & \alpha_! \swarrow & \downarrow k \\ C & \xleftarrow{c_!} & D \end{array}$$

defined as the composition

$$c_!k \xrightarrow{c_!k\eta_a} c_!kaa_! \xrightarrow{c_!\alpha a_!} c_!cha_! \xrightarrow{\varepsilon_c h a_!} ha_!$$

where η_a and ε_c are the unit respectively counit of the adjunctions $a_! \dashv a$ respectively $c_! \dashv c$ fixed above.

The following is a property of the mate that follows easily from the definitions (and actually characterizes it uniquely).

Lemma A.2. *Let the following be a diagram in a 2-category*

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ h \downarrow & \alpha \swarrow & \downarrow k \\ C & \xrightarrow{c} & D \end{array}$$

and $a_! \dashv a$ and $c_! \dashv c$ two fixed adjunctions. Then the following two diagrams commute

$$\begin{array}{ccc} k & \xrightarrow{k\eta_a} & kaa_! \\ \eta_c k \downarrow & & \downarrow \alpha a_! \\ c_!k & \xrightarrow{c_!\alpha_!} & cha_! \end{array} \quad \begin{array}{ccc} c_!ka & \xrightarrow{c_!\alpha_! a} & ha_!a \\ c_!\alpha \downarrow & & \downarrow h\varepsilon_a \\ c_!ch & \xrightarrow{\varepsilon_c h} & h \end{array}$$

where η and ε denote the respective (co)units and $\alpha_!$ is the mate of α .

Proof. This is (the dual of) [Ayo, Proposition 1.1.9]. \square

The following two lemmas express a certain functoriality of the mate construction with respect to pasting of squares. A more abstract (and maybe conceptual) way to formulate them is to present the mate construction as an isomorphism of certain double categories. This can be found in [KS, Proposition 2.2].

Lemma A.3 (Pasting law I). *Let the following be a diagram in a 2-category and its paste*

$$\begin{array}{ccc}
 A & \xrightarrow{a} & B & \xrightarrow{b} & E \\
 \downarrow h & \swarrow \alpha & \downarrow k & \swarrow \beta & \downarrow l \\
 C & \xrightarrow{c} & D & \xrightarrow{d} & F
 \end{array}
 \qquad
 \begin{array}{ccc}
 A & \xrightarrow{ba} & E \\
 \downarrow h & \swarrow \alpha * \beta & \downarrow l \\
 C & \xrightarrow{dc} & F
 \end{array}$$

and $a_! \dashv a$, $b_! \dashv b$, $c_! \dashv c$, and $d_! \dashv d$ four fixed adjunctions. We obtain mates $\alpha_!$ and $\beta_!$ that fit into diagrams of the form

$$\begin{array}{ccc}
 A & \xleftarrow{a_!} & B & \xleftarrow{b_!} & E \\
 \downarrow h & \swarrow \alpha_! & \downarrow k & \swarrow \beta_! & \downarrow l \\
 C & \xleftarrow{c_!} & D & \xleftarrow{d_!} & F
 \end{array}
 \qquad
 \begin{array}{ccc}
 A & \xleftarrow{a_! b_!} & E \\
 \downarrow h & \swarrow \alpha_! * \beta_! & \downarrow l \\
 C & \xleftarrow{c_! d_!} & F
 \end{array}$$

and it holds that $\alpha_! * \beta_! = (\alpha * \beta)_!$, where, for the latter mate, we use the adjunctions $a_! b_! \dashv ba$ and $c_! d_! \dashv dc$ given by composing the original ones.

Proof. This is (the dual of) [Ayo, Proposition 1.1.11] (though note that the composition $\alpha_! * \beta_!$ is erroneously written the wrong way around there, and that what is actually proven is the dual version we stated). \square

Lemma A.4 (Pasting law II). *Let the following be a diagram in a 2-category and its paste*

$$\begin{array}{ccc}
 A & \xrightarrow{a} & B & \xrightarrow{b} & E \\
 \downarrow h & \swarrow \alpha & \downarrow k & \swarrow \beta & \downarrow l \\
 C & \xrightarrow{c} & D & \xrightarrow{d} & F
 \end{array}
 \qquad
 \begin{array}{ccc}
 A & \xrightarrow{ba} & E \\
 \downarrow h & \swarrow \beta * \alpha & \downarrow l \\
 C & \xrightarrow{dc} & F
 \end{array}$$

and $h_! \dashv h$, $k_! \dashv k$, and $l_! \dashv l$ three fixed adjunctions. We obtain mates $\alpha_!$ and $\beta_!$ that fit into diagrams of the form

$$\begin{array}{ccc}
 A & \xrightarrow{a} & B & \xrightarrow{b} & E \\
 \uparrow h_! & \swarrow \alpha_! & \uparrow k_! & \swarrow \beta_! & \uparrow l_! \\
 C & \xrightarrow{c} & D & \xrightarrow{d} & F
 \end{array}
 \qquad
 \begin{array}{ccc}
 A & \xrightarrow{ba} & E \\
 \uparrow h_! & \swarrow \alpha_! * \beta_! & \uparrow l_! \\
 C & \xrightarrow{dc} & F
 \end{array}$$

and it holds that $\alpha_! * \beta_! = (\beta * \alpha)_!$.

Proof. This is (the dual of) [Ayo, Proposition 1.1.12]. \square

Lemma A.5. *Let the following be a diagram in a 2-category*

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ h \downarrow & \alpha \swarrow & \downarrow k \\ C & \xrightarrow{c} & D \end{array}$$

and $\alpha_! \dashv a$ and $c_! \dashv c$ two fixed adjunctions. Furthermore assume that h and k are isomorphisms, and that α is a 2-isomorphism. Then the mate $\alpha_!$ is a 2-isomorphism.

Proof. First note that if $a = c$ (with the same fixed adjunction) and h, k , and α are all identities, then the mate $\alpha_!$ is the identity (by one of the triangle identities). For the general case consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ h \downarrow & \alpha \swarrow & \downarrow k \\ C & \xrightarrow{c} & D \\ h^{-1} \downarrow & \beta \swarrow & \downarrow k^{-1} \\ A & \xrightarrow{a} & B \end{array}$$

where β is given by

$$k^{-1}c = k^{-1}chh^{-1} \xrightarrow{k^{-1}\alpha^{-1}h^{-1}} k^{-1}kah^{-1} = ah^{-1}$$

and note that the paste $\beta * \alpha$ is the identity. This follows from the diagram

$$\begin{array}{ccccc} k^{-1}ka & \xrightarrow{k^{-1}\alpha} & k^{-1}ch & \xrightarrow{\beta h} & ah^{-1}h \\ \parallel & & \parallel & & \parallel \\ k^{-1}kah^{-1}h & \xrightarrow{k^{-1}\alpha h^{-1}h} & k^{-1}chh^{-1}h & \xrightarrow{k^{-1}\alpha^{-1}h^{-1}h} & k^{-1}kah^{-1}h \\ \parallel & & \text{id} & & \parallel \\ a & \xrightarrow{\quad\quad\quad} & & \xrightarrow{\quad\quad\quad} & a \end{array}$$

being commutative. Thus, by the pasting law for mates, we obtain that $h^{-1}\alpha_!$ has a right inverse (namely $\beta_!k$). In the same way we can show that $\alpha_!k^{-1}$ has a left inverse. Since h^{-1} and k^{-1} are both isomorphisms, this implies that $\alpha_!$ has both a left and a right inverse and thus is a 2-isomorphism. \square

Remark A.6. The statement of Lemma A.5 is still true when we only require h and k to be equivalences. Moreover the converse is also true, i.e. if $\alpha_!$ is a 2-isomorphism, then α is as well. (See [GPS, Appendix A].)

Remark A.7. Naturally, there is also a dual version of everything we have done here, using right adjoints instead of left adjoints.

B. Basic ∞ -categorical facts

This appendix consists of a collection of basic ∞ -categorical facts that are used throughout this thesis. They are stated here without proofs, to make it easier to quickly become familiar with the statements. The proofs (or references) can be found in Appendix C. Note that, even though we often only state things for indexing categories (instead of ∞ -categories or simplicial sets), this is purely for convenience and there are more general versions of all of these statements.

Lemma B.1. *Let $f: I \rightarrow J$ and $g: J \rightarrow K$ be maps of simplicial sets and \mathcal{C} an ∞ -category that is both weakly left f -extensible and weakly left g -extensible. Then it is also weakly left $(g \circ f)$ -extensible, and we have $\text{Lan}_{g \circ f} \simeq \text{Lan}_g \circ \text{Lan}_f$.*

Lemma B.2. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a fully faithful functor between categories and \mathcal{C} a left f -extensible ∞ -category. Then the unit $\text{id} \rightarrow \text{Res}_f \text{Lan}_f$ of the adjunction $\text{Lan}_f \dashv \text{Res}_f$ is an equivalence of functors $\text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$.*

Lemma B.3. *Let \mathcal{I} be a category and \mathcal{C} an ∞ -category. Then \mathcal{C} admits all colimits indexed by \mathcal{I} if and only if, for all diagrams $D: \mathcal{I} \rightarrow \mathcal{C}$, there is a colimit diagram extending D . In this case a diagram $D: \mathcal{I}^\triangleright \rightarrow \mathcal{C}$ lies in the essential image of $\text{Lan}_{\text{inc}}: \text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C})$ if and only if it is a colimit diagram.*

Lemma B.4. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories.*

- a) *If \mathcal{J} has a terminal object, then it is contractible.*
- b) *The functor f is homotopy terminal if and only if, for each $j \in \mathcal{J}$, the category $j \downarrow f$ is contractible.*
- c) *If f is right adjoint, then it is homotopy terminal.*
- d) *If f is homotopy terminal, then it is a homotopy equivalence.*

Lemma B.5. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a homotopy terminal functor between categories and \mathcal{C} an ∞ -category that admits colimits indexed both by \mathcal{I} and by \mathcal{J} . Then the natural transformation $f_*: \text{colim}_{\mathcal{I}} \text{Res}_f \rightarrow \text{colim}_{\mathcal{J}}$ of functors $\text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$ is an equivalence.*

Lemma B.6. *Let \mathcal{I} be a category and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor between ∞ -categories that admit colimits indexed by \mathcal{I} . Then the following conditions are equivalent:*

- a) *F preserves left Kan extension along the inclusion $\text{inc}: \mathcal{I} \rightarrow \mathcal{I}^\triangleright$.*
- b) *F preserves colimits indexed by \mathcal{I} .*
- c) *F sends $\mathcal{I}^\triangleright$ -indexed colimit diagrams to colimit diagrams.*

Lemma B.7. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories, $g: \mathcal{K} \rightarrow \mathcal{L}$ a functor between ∞ -categories, and \mathcal{C} a left f -extensible ∞ -category. Then $\text{Fun}(\mathcal{L}, \mathcal{C})$ is left f -extensible, and $\text{Res}_g: \text{Fun}(\mathcal{L}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{K}, \mathcal{C})$ preserves left Kan extension along f .*

C. Tools for Kan extensions and (co)limits

In this appendix we collect some basic tools for working with Kan extensions and (co)limits in ∞ -categories that we need in the rest of this thesis. Note that, even though we often only state things for indexing categories (instead of ∞ -categories or simplicial sets), this is purely for convenience and there are more general versions of most of those statements. Generally, if there is a pair of dual statements, we will only give one of them and leave the other implicit.

There is no claim of originality for any of the statements found in this appendix (the correctness of most, if not all, of them should be more or less clear to anyone familiar with the theory); the ones for which a proof is given are merely those for which the author could not find a good reference.

C.1. Kan extensions

Lemma B.1. *Let $f: I \rightarrow J$ and $g: J \rightarrow K$ be maps of simplicial sets and \mathcal{C} an ∞ -category that is both weakly left f -extensible and weakly left g -extensible. Then it is also weakly left $(g \circ f)$ -extensible, and we have $\text{Lan}_{g \circ f} \simeq \text{Lan}_g \circ \text{Lan}_f$.*

Proof. Since adjunctions compose, we have that $\text{Lan}_g \circ \text{Lan}_f$ is a left adjoint of the composition $\text{Res}_f \circ \text{Res}_g = \text{Res}_{g \circ f}$. As adjoints are unique up to isomorphism (in the homotopy 2-category of ∞ -categories), we obtain that $\text{Lan}_g \circ \text{Lan}_f \simeq \text{Lan}_{g \circ f}$. \square

Lemma B.2. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a fully faithful functor between categories and \mathcal{C} a left f -extensible ∞ -category. Then the unit $\text{id} \rightarrow \text{Res}_f \text{Lan}_f$ of the adjunction $\text{Lan}_f \dashv \text{Res}_f$ is an equivalence of functors $\text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$.*

Proof. This follows from the Beck-Chevalley condition [RV, Proposition 13.5.8], using that by [RV, Lemma 13.4.3] when f is fully faithful a certain square fulfills a condition called exact (here we use that the nerve functor is cosmological by [RV, Example 14.1.1], hence preserves fully faithfulness (cf. [RV, Corollary 3.5.6]) since it preserves absolute right and left lifting diagrams by [RV, Proposition 14.1.9]). \square

Lemma C.1. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a fully faithful functor between categories and \mathcal{C} a left f -extensible ∞ -category. Then $\varepsilon \circ \text{Lan}_f: \text{Lan}_f \text{Res}_f \text{Lan}_f \rightarrow \text{Lan}_f$ is an equivalence, where ε is the counit of the adjunction $\text{Lan}_f \dashv \text{Res}_f$.*

Proof. Consider the following diagram

$$\begin{array}{ccc} \text{Lan}_f & \xrightarrow{\text{id}} & \text{Lan}_f \\ \eta \downarrow & & \uparrow \varepsilon \\ \text{Lan}_f(\text{Res}_f \text{Lan}_f) & \xlongequal{\quad} & (\text{Lan}_f \text{Res}_f) \text{Lan}_f \end{array}$$

where the vertical maps are given by the unit respectively counit of the adjunction $\text{Lan}_f \dashv \text{Res}_f$. It commutes up to homotopy by one of the triangle identities. Since f is fully faithful, the left vertical morphism is an equivalence. Hence the right vertical map is an equivalence as well. \square

Lemma C.2. *Let J be a simplicial set, $f: I \rightarrow I'$ a map of simplicial sets, and \mathcal{C} a weakly left f -extensible ∞ -category. Then there is a homotopy commutative diagram of the form*

$$\begin{array}{ccc}
\mathrm{Fun}(I, \mathrm{Fun}(J, \mathcal{C})) & \xrightarrow{\mathrm{Lan}_f} & \mathrm{Fun}(I', \mathrm{Fun}(J, \mathcal{C})) \\
\cong \uparrow & & \uparrow \cong \\
\mathrm{Fun}(I \times J, \mathcal{C}) & \xrightarrow{\mathrm{Lan}_{f \times \mathrm{id}}} & \mathrm{Fun}(I' \times J, \mathcal{C}) \\
\cong \downarrow & & \downarrow \cong \\
\mathrm{Fun}(J, \mathrm{Fun}(I, \mathcal{C})) & \xrightarrow{\mathrm{Lan}_f \circ} & \mathrm{Fun}(J, \mathrm{Fun}(I', \mathcal{C}))
\end{array}$$

where the vertical maps are the respective currying isomorphisms (in particular, all of these left Kan extensions actually exist).

Proof. Note that $(\mathrm{Lan}_f \circ)$ is left adjoint to $(\mathrm{Res}_f \circ)$ by Lemma D.5. Since the above diagram with the restrictions, instead of their left adjoints, commutes, we obtain that Res_f and $\mathrm{Res}_{f \times \mathrm{id}}$ actually have left adjoints. Then Lemma A.5 implies the statement. \square

Lemma C.3. *Let I and J be simplicial sets, \mathcal{C} an ∞ -category that admits colimits indexed by I , and $D: I \times J \rightarrow \mathcal{C}$ a functor. Denote by $D_I: I \rightarrow \mathrm{Fun}(J, \mathcal{C})$ and $D_J: J \rightarrow \mathrm{Fun}(I, \mathcal{C})$ the curried functors. Then $\mathrm{Lan}_{\mathrm{pr}_J} D$, $\mathrm{colim}_I D_I$, and $\mathrm{colim}_I \circ D_J$ exist and are all equivalent in $\mathrm{Fun}(J, \mathcal{C})$.*

Proof. This is a special case of Lemma C.2. \square

Lemma C.4. *Let $f: \mathcal{I} \rightarrow \mathcal{K}$ and $g: \mathcal{J} \rightarrow \mathcal{K}$ be functors between categories. Consider the natural transformation*

$$\begin{array}{ccc}
f \downarrow g & \xrightarrow{\mathrm{pr}_{\mathcal{I}}} & \mathcal{I} \\
\mathrm{pr}_{\mathcal{J}} \downarrow & \swarrow \alpha & \downarrow f \\
\mathcal{J} & \xrightarrow{g} & \mathcal{K}
\end{array}$$

given, at $(i, j, k: f(i) \rightarrow g(j)) \in f \downarrow g$, by k . Now let \mathcal{C} be a left f -extensible and left $\mathrm{pr}_{\mathcal{J}}$ -extensible ∞ -category. After applying $\mathrm{Fun}(-, \mathcal{C})$ to the diagram above we get

$$\begin{array}{ccc}
\mathrm{Fun}(\mathcal{K}, \mathcal{C}) & \xrightarrow{\mathrm{Res}_f} & \mathrm{Fun}(\mathcal{I}, \mathcal{C}) \\
\mathrm{Res}_g \downarrow & \swarrow \alpha & \downarrow \mathrm{Res}_{\mathrm{pr}_{\mathcal{I}}} \\
\mathrm{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\mathrm{Res}_{\mathrm{pr}_{\mathcal{J}}}} & \mathrm{Fun}(f \downarrow g, \mathcal{C})
\end{array}$$

and taking the mate we obtain a transformation $\alpha_1: \mathrm{Lan}_{\mathrm{pr}_{\mathcal{J}}} \mathrm{Res}_{\mathrm{pr}_{\mathcal{I}}} \rightarrow \mathrm{Res}_g \mathrm{Lan}_f$. This transformation α_1 is an equivalence.

Proof. This follows from the fact that the second diagram satisfies the Beck-Chevalley condition by [RV, Proposition 13.5.8] as the first one is a so called exact square by [RV, Lemma 13.2.5] (again using that the nerve is a cosmological functor by [RV, Example 14.1.1] and thus preserves comma categories by [RV, Proposition 14.1.7]). \square

Lemma C.5. *Let \mathcal{I} and \mathcal{J} be categories, $f: \mathcal{I} \rightarrow \mathcal{J}$ a functor, \mathcal{C} a left f -extensible ∞ -category, and $D: \mathcal{I} \rightarrow \mathcal{C}$ a diagram. Then, for any $j \in \mathcal{J}$, the mate $\vartheta: \operatorname{colim}_{f \downarrow j} \operatorname{Res}_{\operatorname{pr}} \rightarrow \operatorname{Res}_j \operatorname{Lan}_f$ of the natural transformation in the right diagram (which is the image of the left diagram under $\operatorname{Fun}(-, \mathcal{C})$)*

$$\begin{array}{ccc} f \downarrow j & \xrightarrow{\operatorname{const}} & * \\ \operatorname{pr} \downarrow & \nearrow \tilde{\rho}_j & \downarrow j \\ \mathcal{I} & \xrightarrow{f} & \mathcal{J} \end{array} \qquad \begin{array}{ccc} \operatorname{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\operatorname{Res}_f} & \operatorname{Fun}(\mathcal{I}, \mathcal{C}) \\ \operatorname{Res}_j \downarrow & \nearrow \rho_j & \downarrow \operatorname{Res}_{\operatorname{pr}} \\ \mathcal{C} & \xrightarrow{\Delta} & \operatorname{Fun}(f \downarrow j, \mathcal{C}) \end{array}$$

is an equivalence, where $\tilde{\rho}_j$ is, at $(i, k: f(i) \rightarrow j)$, just given by k . Furthermore it is natural in j , in the sense that, for a morphism $\kappa: j \rightarrow j'$ in \mathcal{J} , the diagram

$$\begin{array}{ccc} \operatorname{colim}_{f \downarrow j} \operatorname{Res}_{\operatorname{pr}} & \xrightarrow{(f \downarrow \kappa)_*} & \operatorname{colim}_{f \downarrow j'} \operatorname{Res}_{\operatorname{pr}} \\ \vartheta \downarrow \simeq & & \simeq \downarrow \vartheta \\ \operatorname{Res}_j \operatorname{Lan}_f & \longrightarrow & \operatorname{Res}_{j'} \operatorname{Lan}_f \end{array}$$

commutes up to homotopy.

Proof. That ϑ is an equivalence is a special case of Lemma C.4. For the naturality in j we consider, for a map $\kappa: j \rightarrow j'$, the two diagrams

$$\begin{array}{ccc} f \downarrow j & \xrightarrow{f \downarrow \kappa} & f \downarrow j' & \xrightarrow{\operatorname{pr}} & \mathcal{I} \\ \downarrow & \swarrow \operatorname{id} & \downarrow & \swarrow \tilde{\rho}_{j'} & \downarrow f \\ * & \longrightarrow & * & \longrightarrow & \mathcal{J} \end{array} \qquad \begin{array}{ccc} f \downarrow j & \longrightarrow & * & \longrightarrow & * \\ \operatorname{pr} \downarrow & \nearrow \tilde{\rho}_j & \downarrow j & \nearrow \kappa & \downarrow j' \\ \mathcal{I} & \xrightarrow{f} & \mathcal{J} & \xrightarrow{\operatorname{id}} & \mathcal{J} \end{array}$$

for which we note that $\operatorname{id} * \tilde{\rho}_{j'} = \kappa * \tilde{\rho}_j$ by definition of the involved maps. Hence, after applying $\operatorname{Fun}(-, \mathcal{C})$, we obtain, by the pasting laws for mates, that $(\rho_{j'})_! * \operatorname{id}_! = (\operatorname{id} * \rho_{j'})_! = (\kappa * \rho_j)_! = \kappa_! * (\rho_j)_!$ (in the homotopy 2-category of ∞ -categories). This is the statement we wanted to show since $\kappa_!$ is the map $\operatorname{Res}_j \rightarrow \operatorname{Res}_{j'}$ given by evaluation at κ and $\operatorname{id}_!$ is the map on colimits induced by $f \downarrow \kappa$. \square

Lemma C.6. *Let the diagram in the left be a diagram of categories, functors between them, and a natural transformation and the one in the right its image under $\operatorname{Fun}(-, \mathcal{C})$*

$$\begin{array}{ccc} \mathcal{I} & \xrightarrow{a} & \mathcal{J} \\ b \downarrow & \nearrow \gamma & \downarrow d \\ \mathcal{K} & \xrightarrow{c} & \mathcal{L} \end{array} \qquad \begin{array}{ccc} \operatorname{Fun}(\mathcal{L}, \mathcal{C}) & \xrightarrow{\operatorname{Res}_d} & \operatorname{Fun}(\mathcal{J}, \mathcal{C}) \\ \operatorname{Res}_c \downarrow & \nearrow \gamma & \downarrow \operatorname{Res}_a \\ \operatorname{Fun}(\mathcal{K}, \mathcal{C}) & \xrightarrow{\operatorname{Res}_b} & \operatorname{Fun}(\mathcal{I}, \mathcal{C}) \end{array}$$

where \mathcal{C} is a left b -extensible and left d -extensible ∞ -category. Then, for any $k \in \mathcal{K}$, the following diagram commutes up to homotopy

$$\begin{array}{ccc} \operatorname{colim}_{b \downarrow k} \operatorname{Res}_{\operatorname{pr}_{b \downarrow k}} \operatorname{Res}_a & \xrightarrow{f_*} & \operatorname{colim}_{d \downarrow c(k)} \operatorname{Res}_{\operatorname{pr}_{d \downarrow c(k)}} \\ \vartheta \downarrow \simeq & & \simeq \downarrow \vartheta \\ \operatorname{Res}_k \operatorname{Lan}_b \operatorname{Res}_a & \xrightarrow{\gamma!} & \operatorname{Res}_k \operatorname{Res}_c \operatorname{Lan}_d \end{array}$$

where $\gamma!$ is the mate of γ , the maps denoted ϑ are as in Lemma C.5, and f is the functor

$$b \downarrow k \longrightarrow d \downarrow c(k), \quad (i, b(i) \xrightarrow{g} k) \longmapsto (a(i), d(a(i)) \xrightarrow{\gamma} c(b(i)) \xrightarrow{c(g)} c(k))$$

acting on morphisms via a .

Proof. Consider the two diagrams

$$\begin{array}{ccc} b \downarrow k & \xrightarrow{\operatorname{pr}_{b \downarrow k}} & \mathcal{I} & \xrightarrow{a} & \mathcal{J} \\ \downarrow & \swarrow \tilde{\rho}_k & \downarrow b & \swarrow \gamma & \downarrow d \\ * & \xrightarrow{k} & \mathcal{K} & \xrightarrow{c} & \mathcal{L} \end{array} \qquad \begin{array}{ccc} b \downarrow k & \xrightarrow{f} & d \downarrow c(k) & \xrightarrow{\operatorname{pr}_{d \downarrow c(k)}} & \mathcal{J} \\ \downarrow & \swarrow \operatorname{id} & \downarrow & \swarrow \tilde{\rho}_{c(k)} & \downarrow d \\ * & \longrightarrow & * & \xrightarrow{c(k)} & \mathcal{L} \end{array}$$

where $\tilde{\rho}$ is as in Lemma C.5. Note that it follows directly from the definitions that their pastes $\tilde{\rho}_k * \gamma$ and $\operatorname{id} * \tilde{\rho}_{c(k)}$ are the same. Applying $\operatorname{Fun}(-, \mathcal{C})$ and using the pasting law for mates yields the desired result. \square

C.2. (Co)Limits

Lemma C.7. *Let \mathcal{C} be an ∞ -category that admits colimits indexed by a category \mathcal{I} . Then the mate $\operatorname{colim}_{\mathcal{I}} \rightarrow \operatorname{Res}_{\blacktriangleright} \operatorname{Lan}_{\operatorname{inc}}$ of the diagram on the right (which is the image of the diagram on the left under $\operatorname{Fun}(-, \mathcal{C})$)*

$$\begin{array}{ccc} \mathcal{I} & \xrightarrow{\operatorname{id}} & \mathcal{I} \\ \downarrow & \swarrow \alpha & \downarrow \operatorname{inc} \\ * & \xrightarrow{\blacktriangleright} & \mathcal{I}^{\blacktriangleright} \end{array} \qquad \begin{array}{ccc} \operatorname{Fun}(\mathcal{I}^{\blacktriangleright}, \mathcal{C}) & \xrightarrow{\operatorname{Res}_{\operatorname{inc}}} & \operatorname{Fun}(\mathcal{I}, \mathcal{C}) \\ \operatorname{Res}_{\blacktriangleright} \downarrow & \swarrow \alpha & \downarrow \operatorname{id} \\ \mathcal{C} & \xrightarrow{\Delta} & \operatorname{Fun}(\mathcal{I}, \mathcal{C}) \end{array}$$

is an equivalence.

Proof. This is a special case of Lemma C.5 since there is an isomorphism $\operatorname{inc} \downarrow \blacktriangleright \cong \mathcal{I}$ over $\mathcal{I}^{\blacktriangleright}$. \square

Lemma B.3. *Let \mathcal{I} be a category and \mathcal{C} an ∞ -category. Then \mathcal{C} admits all colimits indexed by \mathcal{I} if and only if, for all diagrams $D: \mathcal{I} \rightarrow \mathcal{C}$, there is a colimit diagram extending D . In this case a diagram $D: \mathcal{I}^{\blacktriangleright} \rightarrow \mathcal{C}$ lies in the essential image of $\operatorname{Lan}_{\operatorname{inc}}: \operatorname{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \operatorname{Fun}(\mathcal{I}^{\blacktriangleright}, \mathcal{C})$ if and only if it is a colimit diagram.*

Proof. The first statement follows from [RV, Proposition F.2.1 and Corollary 16.2.10]. The second from [RV, Proposition F.2.1, Lemma 2.3.5, and Lemma 2.3.6]. \square

Lemma B.4. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories.*

- a) *If \mathcal{J} has a terminal object, then it is contractible.*
- b) *The functor f is homotopy terminal if and only if, for each $j \in \mathcal{J}$, the category $j \downarrow f$ is contractible.*
- c) *If f is right adjoint, then it is homotopy terminal.*
- d) *If f is homotopy terminal, then it is a homotopy equivalence.*

Proof. The first statement is clear (one can explicitly construct the contraction). The latter three statements follow, in order, from [LurHTT, Theorem 4.1.3.1], [RV, Proposition 4.1.4], and [LurHTT, Proposition 4.1.1.3]. \square

Lemma C.8. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a homotopy terminal functor between categories and \mathcal{C} an ∞ -category. Then \mathcal{C} admits colimits indexed by \mathcal{I} if and only if it admits colimits indexed by \mathcal{J} .*

Proof. This follows from Lemma B.3 and [LurHTT, Proposition 4.1.1.8 (2)]. \square

Lemma B.5. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a homotopy terminal functor between categories and \mathcal{C} an ∞ -category that admits colimits indexed both by \mathcal{I} and by \mathcal{J} . Then the natural transformation $f_*: \text{colim}_{\mathcal{I}} \text{Res}_f \rightarrow \text{colim}_{\mathcal{J}}$ of functors $\text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$ is an equivalence.*

Proof. By Lemma C.6, the mate $\text{id}_! : \text{Lan}_{\text{inc}_{\mathcal{I}}} \text{Res}_f \rightarrow \text{Res}_{f \triangleright} \text{Lan}_{\text{inc}_{\mathcal{J}}}$ of the natural transformation in the diagram on the right (which is the image of the diagram on the left under $\text{Fun}(-, \mathcal{C})$)

$$\begin{array}{ccc}
 \mathcal{I} & \xrightarrow{f} & \mathcal{J} \\
 \text{inc} \downarrow & \text{id} \swarrow & \downarrow \text{inc} \\
 \mathcal{I}^{\triangleright} & \xrightarrow{f^{\triangleright}} & \mathcal{J}^{\triangleright}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{Fun}(\mathcal{J}^{\triangleright}, \mathcal{C}) & \xrightarrow{\text{Res}_{\text{inc}}} & \text{Fun}(\mathcal{J}, \mathcal{C}) \\
 \text{Res}_{f^{\triangleright}} \downarrow & \text{id} \swarrow & \downarrow \text{Res}_f \\
 \text{Fun}(\mathcal{I}^{\triangleright}, \mathcal{C}) & \xrightarrow{\text{Res}_{\text{inc}}} & \text{Fun}(\mathcal{I}, \mathcal{C})
 \end{array}$$

is given, at the cocone point, by f_* . Hence it is enough to prove that $\text{id}_!$ is an equivalence. This mate is given by the composition

$$\begin{array}{ccc}
 \text{Lan}_{\text{inc}_{\mathcal{I}}} \text{Res}_f & \xrightarrow{\eta} & \text{Lan}_{\text{inc}_{\mathcal{I}}} \text{Res}_f \text{Res}_{\text{inc}_{\mathcal{J}}} \text{Lan}_{\text{inc}_{\mathcal{J}}} \\
 & & \parallel \\
 & & \text{Lan}_{\text{inc}_{\mathcal{I}}} \text{Res}_{\text{inc}_{\mathcal{I}}} \text{Res}_{f^{\triangleright}} \text{Lan}_{\text{inc}_{\mathcal{J}}} \xrightarrow{\varepsilon} \text{Res}_{f^{\triangleright}} \text{Lan}_{\text{inc}_{\mathcal{J}}}
 \end{array}$$

of the unit η of the adjunction $\text{Lan}_{\text{inc}_{\mathcal{J}}} \dashv \text{Res}_{\text{inc}_{\mathcal{J}}}$ and the counit ε of the adjunction $\text{Lan}_{\text{inc}_{\mathcal{I}}} \dashv \text{Res}_{\text{inc}_{\mathcal{I}}}$. Since $\text{inc}_{\mathcal{J}}$ is fully faithful, the map η is an equivalence. So we only need to show that $\varepsilon \circ \text{Res}_{f^{\triangleright}} \circ \text{Lan}_{\text{inc}_{\mathcal{J}}}$ is an equivalence. Let $D: \mathcal{I} \rightarrow \mathcal{C}$ be a diagram. By Lemma B.3 and assumption the diagram $(\text{Res}_{f^{\triangleright}} \circ \text{Lan}_{\text{inc}_{\mathcal{J}}})(D)$ is a colimit diagram. But ε applied to a colimit diagram is an equivalence by Lemma C.1 and again Lemma B.3. \square

Lemma C.9. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories and \mathcal{C} an ∞ -category that admits colimits indexed both by \mathcal{I} and by \mathcal{J} . Then the following diagram in $\text{Fun}(\text{Fun}(\mathcal{J}^\triangleright, \mathcal{C}), \mathcal{C})$ commutes up to homotopy*

$$\begin{array}{ccc}
\text{colim}_{\mathcal{I}} \text{Res}_f \text{Res}_{\text{inc}_{\mathcal{J}}} & \xrightarrow{f_*} & \text{colim}_{\mathcal{J}} \text{Res}_{\text{inc}_{\mathcal{J}}} \\
\parallel & & \downarrow \\
\text{colim}_{\mathcal{I}} \text{Res}_{\text{inc}_{\mathcal{I}}} \text{Res}_{f^\triangleright} & & \\
\downarrow & & \downarrow \\
\text{Res}_{\blacktriangleright} \text{Res}_{f^\triangleright} & \xlongequal{\quad} & \text{Res}_{\blacktriangleright}
\end{array}$$

where the vertical maps are the canonical maps out of the colimit.

Proof. Consider the two diagrams

$$\begin{array}{ccc}
\mathcal{I} & \xrightarrow{\text{inc}} & \mathcal{I}^\triangleright & \xrightarrow{f^\triangleright} & \mathcal{J}^\triangleright \\
\downarrow & \xi_{\mathcal{I}} \swarrow & \text{id} \downarrow & \text{id}_1 \swarrow & \downarrow \text{id} \\
* & \xrightarrow{\blacktriangleright} & \mathcal{I}^\triangleright & \xrightarrow{f^\triangleright} & \mathcal{J}^\triangleright
\end{array}
\qquad
\begin{array}{ccc}
\mathcal{I} & \xrightarrow{f} & \mathcal{J} & \xrightarrow{\text{inc}} & \mathcal{J}^\triangleright \\
\downarrow & \text{id}_2 \swarrow & \downarrow & \xi_{\mathcal{J}} \swarrow & \downarrow \text{id} \\
* & \xrightarrow{\blacktriangleright} & * & \xrightarrow{\blacktriangleright} & \mathcal{J}^\triangleright
\end{array}$$

where $\xi_{\mathcal{I}}$ and $\xi_{\mathcal{J}}$ are as in Notation 1.25, and note that their pastes agree, i.e. $\xi_{\mathcal{I}} * \text{id}_1 = \text{id}_2 * \xi_{\mathcal{J}}$. Applying $\text{Fun}(-, \mathcal{C})$ and using the pasting law for mates yields $(\text{id}_1)_! * (\xi_{\mathcal{I}})_! \simeq (\xi_{\mathcal{J}})_! * (\text{id}_2)_!$. This is what we wanted to show since $(\text{id}_2)_! = f_*$ and $(\text{id}_1)_!$ is the identity. \square

Lemma C.10. *Let \mathcal{I} be a category with a terminal object $*$, \mathcal{C} an ∞ -category, and $D: \mathcal{I}^\triangleright \rightarrow \mathcal{C}$ a diagram such that D applied to the unique morphism $* \rightarrow \blacktriangleright$ is an equivalence. Then the canonical morphism $\text{colim}_{\mathcal{I}} D|_{\mathcal{I}} \rightarrow D(\blacktriangleright)$ is an equivalence as well.*

Proof. First note that \mathcal{C} admits colimits indexed by \mathcal{I} by Lemma C.8. Applying Lemma C.9 to the functor $\text{const}_*: * \rightarrow \mathcal{I}$, we obtain a diagram

$$\begin{array}{ccc}
\text{colim}_* D(*) & \xrightarrow{\simeq} & \text{colim}_{\mathcal{I}} D|_{\mathcal{I}} \\
\downarrow \simeq & & \downarrow \\
D(\blacktriangleright) & \xlongequal{\quad} & D(\blacktriangleright)
\end{array}$$

where the top horizontal morphism is an equivalence since const_* is homotopy terminal, and the left vertical morphism is an equivalence by Remark 1.26. \square

Lemma C.11. *Let \mathcal{I} be a category, \mathcal{C} an ∞ -category that admits colimits indexed by \mathcal{I} , and $D: \mathcal{I}^\triangleright \rightarrow \mathcal{C}$ a diagram. Then the canonical map $(\text{colim}_{\mathcal{I}} \text{Res}_{\text{inc}})(D) \rightarrow \text{Res}_{\blacktriangleright}(D)$ is an equivalence if and only if the counit $\text{Lan}_{\text{inc}} \text{Res}_{\text{inc}} \rightarrow \text{id}$ of the adjunction $\text{Lan}_{\text{inc}} \dashv \text{Res}_{\text{inc}}$ is an equivalence at D .*

Proof. The mate of the natural transformation on the right (which is the image under $\text{Fun}(-, \mathcal{C})$ of the natural transformation on the left)

$$\begin{array}{ccc}
\mathcal{I} & \xrightarrow{\text{inc}} & \mathcal{T}^\triangleright \\
\text{inc} \downarrow & \text{id} \swarrow & \downarrow \text{id} \\
\mathcal{T}^\triangleright & \xrightarrow{\text{id}} & \mathcal{T}^\triangleright
\end{array}
\qquad
\begin{array}{ccc}
\text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) & \xrightarrow{\text{id}} & \text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) \\
\text{id} \downarrow & \text{id} \swarrow & \downarrow \text{Res}_{\text{inc}} \\
\text{Fun}(\mathcal{T}^\triangleright, \mathcal{C}) & \xrightarrow{\text{Res}_{\text{inc}}} & \text{Fun}(\mathcal{I}, \mathcal{C})
\end{array}$$

is precisely the counit $\text{Lan}_{\text{inc}} \text{Res}_{\text{inc}} \rightarrow \text{id}$. Hence, by Lemma C.6, it is an equivalence at D if and only if, for all $k \in \mathcal{T}^\triangleright$, the map

$$\text{colim}_{\text{inc} \downarrow k} \text{Res}_{\text{pr}_{\text{inc} \downarrow k}} \text{Res}_{\text{inc}} = \text{colim}_{\text{inc} \downarrow k} \text{Res}_{f_k} \text{Res}_{\text{pr}_{\text{id} \downarrow k}} \xrightarrow{(f_k)_*} \text{colim}_{\text{id} \downarrow k} \text{Res}_{\text{pr}_{\text{id} \downarrow k}}$$

is an equivalence at D , where $f_k: \text{inc} \downarrow k \rightarrow \text{id} \downarrow k$ is the canonical inclusion. When k is not \blacktriangleright , then f_k is an isomorphism and $(f_k)_*$ is an equivalence. When k is \blacktriangleright , then f_k is just (isomorphic to) inc , and $\text{pr}_{\text{id} \downarrow k}$ is an isomorphism. So it is enough to show that $\text{inc}_*: \text{colim}_{\mathcal{I}} \text{Res}_{\text{inc}} \rightarrow \text{colim}_{\mathcal{T}^\triangleright}$ is an equivalence at D if and only if the canonical map $(\text{colim}_{\mathcal{I}} \text{Res}_{\text{inc}})(D) \rightarrow \text{Res}_{\blacktriangleright}(D)$ is an equivalence. This follows from Lemma C.9 by considering the diagram $D': (\mathcal{T}^\triangleright)^\triangleright \rightarrow \mathcal{C}$ obtained from D by pulling back along the functor $(\mathcal{T}^\triangleright)^\triangleright \rightarrow \mathcal{T}^\triangleright$ that is the identity on $\mathcal{T}^\triangleright$ and sends the new cocone point to the old one (here we also use Lemma C.10 to see that canonical map $(\text{colim}_{\mathcal{T}^\triangleright} \text{Res}_{\text{inc}_{\mathcal{T}^\triangleright}})(D') \rightarrow \text{Res}_{\blacktriangleright}(D')$ is an equivalence). \square

Lemma C.12. *Let I and J be simplicial sets, \mathcal{C} an ∞ -category that admits colimits indexed by J , and $f: I \rightarrow \text{Fun}(J^\triangleright, \mathcal{C})$ a functor. Denote by $g: J^\triangleright \rightarrow \text{Fun}(I, \mathcal{C})$ the functor obtained from f via currying. Then there is a homotopy commutative diagram of the form*

$$\begin{array}{ccc}
\text{Res}_{\blacktriangleright} \circ f & \longleftarrow & \text{colim}_J \circ \text{Res}_{\text{inc}} \circ f \\
\parallel & & \downarrow \simeq \\
\text{Res}_{\blacktriangleright} g & \longleftarrow & \text{colim}_J (\text{Res}_{\text{inc}} g)
\end{array} \tag{C.1}$$

where the horizontal morphisms are the canonical maps from the colimit.

Proof. Consider the two diagrams

$$\begin{array}{ccc}
\text{Fun}(J^\triangleright, \text{Fun}(I, \mathcal{C})) & \xrightarrow{\text{id}} & \text{Fun}(J^\triangleright, \text{Fun}(I, \mathcal{C})) \\
\text{Res}_{\blacktriangleright} \downarrow & \text{id}_1 \swarrow & \downarrow \text{Res}_{\text{inc}} \\
\text{Fun}(I, \mathcal{C}) & \xrightarrow{\Delta} & \text{Fun}(J, \text{Fun}(I, \mathcal{C})) \\
\text{id} \downarrow & \text{id}_2 \swarrow & \downarrow \cong \\
\text{Fun}(I, \mathcal{C}) & \xrightarrow{\Delta \circ} & \text{Fun}(I, \text{Fun}(J, \mathcal{C}))
\end{array}$$

and

$$\begin{array}{ccc}
\text{Fun}(J^\triangleright, \text{Fun}(I, \mathcal{C})) & \xrightarrow{\text{id}} & \text{Fun}(J^\triangleright, \text{Fun}(I, \mathcal{C})) \\
\cong \downarrow & \swarrow \text{id}_3 & \downarrow \cong \\
\text{Fun}(I, \text{Fun}(J^\triangleright, \mathcal{C})) & \xrightarrow{\text{id}} & \text{Fun}(I, \text{Fun}(J^\triangleright, \mathcal{C})) \\
\text{Res}_\blacktriangleright \circ \downarrow & \swarrow \text{id}_4 & \downarrow \text{Res}_{\text{inc}} \circ \\
\text{Fun}(I, \mathcal{C}) & \xrightarrow{\Delta \circ} & \text{Fun}(I, \text{Fun}(J, \mathcal{C}))
\end{array}$$

and note that their pastes agree. Now the mate of id_1 gives the lower horizontal map in diagram (C.1), the mate of id_4 the upper horizontal map (where we use the adjunction $(\text{colim}_J \circ) \dashv (\Delta \circ)$ obtained from Lemma D.5), the mate of id_2 the right equivalence (using Lemma A.5), and the mate of id_3 the left identity. An application of the pasting law for mates yields the desired statement. \square

Lemma C.13. *Let \mathcal{I} be a contractible category, \mathcal{C} an ∞ -category that admits colimits indexed by \mathcal{I} , and $D: \mathcal{I} \rightarrow \mathcal{C}$ a diagram such that, for all morphisms k of \mathcal{I} , the induced map $D(k)$ is an equivalence. Then, for all $i \in \mathcal{I}$, the structure map $D(i) \rightarrow \text{colim}_{\mathcal{I}} D$ is an equivalence.*

Proof. We will show that the composition

$$\text{Fun}(\mathcal{I}, \mathcal{C}) \xrightarrow{\text{Lan}_{\text{inc}}} \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C}) \xrightarrow{\text{Res}_{t_i}} \text{Fun}(\Delta^1, \mathcal{C})$$

sends D to an equivalence, where $t_i: \Delta^1 \rightarrow \mathcal{I}^\triangleright$ is as in Notation 1.23, i.e. the functor representing the unique morphism $i \rightarrow \blacktriangleright$. Note that $\text{Lan}_{\text{inc}} D$ is a colimit diagram indexed by $\mathcal{I}^\triangleright$ that sends any morphism in \mathcal{I} to an equivalence. Hence, by [LurHTT, Proposition 4.3.1.12] (together with [LurHTT, Proposition 2.4.1.5]), the diagram $\text{Lan}_{\text{inc}} D$ sends every morphism of $\mathcal{I}^\triangleright$ to an equivalence, which implies the claim. \square

C.3. Preservation of Kan extensions and (co)limits

Lemma B.7. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories, $g: \mathcal{K} \rightarrow \mathcal{L}$ a functor between ∞ -categories, and \mathcal{C} a left f -extensible ∞ -category. Then $\text{Fun}(\mathcal{L}, \mathcal{C})$ is left f -extensible, and $\text{Res}_g: \text{Fun}(\mathcal{L}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{K}, \mathcal{C})$ preserves left Kan extension along f .*

Proof. That $\text{Fun}(\mathcal{L}, \mathcal{C})$ is weakly left f -extensible when \mathcal{C} is so was part of Lemma C.2. This also implies the corresponding statement for left f -extensible since this was defined as certain colimits existing which in turn was defined via weakly const-extensible.

For the second part we want that the mate of the transformation id_2 in the diagram

$$\begin{array}{ccc}
\text{Fun}(\mathcal{J} \times \mathcal{L}, \mathcal{C}) & \xrightarrow{\text{Res}_{f \times \text{id}}} & \text{Fun}(\mathcal{I} \times \mathcal{L}, \mathcal{C}) \\
\downarrow \cong & \swarrow \text{id}_1 & \downarrow \cong \\
\text{Fun}(\mathcal{J}, \text{Fun}(\mathcal{L}, \mathcal{C})) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{L}, \mathcal{C})) \\
\downarrow \text{Res}_g \circ & \swarrow \text{id}_2 & \downarrow \text{Res}_g \circ \\
\text{Fun}(\mathcal{J}, \text{Fun}(\mathcal{K}, \mathcal{D})) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{K}, \mathcal{D})) \\
\downarrow \cong & \swarrow \text{id}_3 & \downarrow \cong \\
\text{Fun}(\mathcal{J} \times \mathcal{K}, \mathcal{D}) & \xrightarrow{\text{Res}_{f \times \text{id}}} & \text{Fun}(\mathcal{I} \times \mathcal{K}, \mathcal{D})
\end{array}$$

$\text{Res}_{\text{id} \times g}$ $\text{Res}_{\text{id} \times g}$

is an equivalence. For this note that the mates of id_1 and id_3 are equivalences by Lemma A.5 and that the paste of all three transformations is just $\text{Fun}(-, \mathcal{C})$ applied to the transformation

$$\begin{array}{ccc}
\mathcal{I} \times \mathcal{K} & \xrightarrow{\text{id} \times g} & \mathcal{I} \times \mathcal{L} \\
f \times \text{id} \downarrow & \swarrow \text{id} & \downarrow f \times \text{id} \\
\mathcal{J} \times \mathcal{K} & \xrightarrow{\text{id} \times g} & \mathcal{J} \times \mathcal{L}
\end{array}$$

which is a so called exact square by [RV, Lemma 13.2.7]. Hence, again by the Beck-Chevalley condition [RV, Proposition 13.5.8] (which we can use since the argument in Remark 1.20 applies to $f \times \text{id}$ as its slice ∞ -categories admit homotopy terminal functors from those of f), the mate of this paste is an equivalence. Now the pasting law for mates implies that the mate of id_2 is an equivalence which we wanted to show. \square

Lemma C.14. *Let \mathcal{K} be an ∞ -category, $f: \mathcal{I} \rightarrow \mathcal{J}$ a functor between categories, \mathcal{C} and \mathcal{D} two left f -extensible ∞ -categories, and $F: \mathcal{C} \rightarrow \text{Fun}(\mathcal{K}, \mathcal{D})$ a functor. Then F preserves left Kan extension along f if and only if, for all $k \in \mathcal{K}$, the functor $\text{Res}_k \circ F: \mathcal{C} \rightarrow \mathcal{D}$ preserves left Kan extension along f .*

Proof. Consider the diagram

$$\begin{array}{ccc}
\text{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
F \circ \downarrow & \swarrow \text{id}_1 & \downarrow F \circ \\
\text{Fun}(\mathcal{J}, \text{Fun}(\mathcal{K}, \mathcal{D})) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{K}, \mathcal{D})) \\
\downarrow \text{Res}_k \circ & \swarrow \text{id}_2 & \downarrow \text{Res}_k \circ \\
\text{Fun}(\mathcal{J}, \mathcal{D}) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \mathcal{D})
\end{array}$$

and note that $(\text{id}_1)_! * (\text{id}_2)_! \simeq (\text{id}_2 * \text{id}_1)_!$ by the pasting law for mates. Since Res_k preserves left Kan extension along f , the mate $(\text{id}_2)_!$ is an equivalence. Now, noting that $(\text{Res}_k \circ) \circ (\text{id}_1)_!$ is the other part of the composition in the paste $(\text{id}_1)_! * (\text{id}_2)_!$, we obtain that $(\text{Res}_k \circ) \circ (\text{id}_1)_!$ is an equivalence if and only if $(\text{id}_2 * \text{id}_1)_!$ is an equivalence. As the former being true for all k is equivalent to F preserving left Kan extension along f , and the latter is the definition of $\text{Res}_k \circ F$ preserving left Kan extension along f , this implies the claim. \square

Lemma C.15. *Let \mathcal{J} and \mathcal{J} be ∞ -categories, \mathcal{K} a category, and \mathcal{C} and \mathcal{D} two ∞ -categories that admit colimits indexed by \mathcal{K} .*

- a) *Let $g: \mathcal{C} \rightarrow \mathcal{D}$ be a functor that preserves colimits indexed by \mathcal{K} . Then the induced functor $(g \circ): \text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{J}, \mathcal{D})$ preserves colimits indexed by \mathcal{K} .*
- b) *The functor $h: \text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \text{Fun}(\text{Fun}(\mathcal{J}, \mathcal{J}), \text{Fun}(\mathcal{J}, \mathcal{C}))$ given by $f \mapsto (f \circ)$ preserves colimits indexed by \mathcal{K} .*

Proof. The first statement follows from Lemma C.14 since, for all $i \in \mathcal{J}$, it holds that $\text{Res}_i \circ (g \circ) = g \circ \text{Res}_i$ and both functors in the latter composition preserve colimits indexed by \mathcal{K} . The second statement follows from the same lemma by noting that, for any $f \in \text{Fun}(\mathcal{J}, \mathcal{J})$, the functor $\text{Res}_{\{f\}} \circ h = \text{Res}_f$ preserves colimits indexed by \mathcal{K} . \square

Lemma C.16. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ be a functor between categories, \mathcal{C} and \mathcal{D} two left f -extensible ∞ -categories, and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor that preserves colimits indexed by $f \downarrow j$ for all $j \in \mathcal{J}$. Then F preserves left Kan extensions along f .*

Proof. Consider the two diagrams

$$\begin{array}{ccc}
 \text{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
 F \circ \downarrow & \text{id}_1 \swarrow & \downarrow F \circ \\
 \text{Fun}(\mathcal{J}, \mathcal{D}) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \mathcal{D}) \\
 \text{Res}_j \downarrow & \rho_j \swarrow & \downarrow \text{Res}_{\text{pr}} \\
 \mathcal{D} & \xrightarrow{\Delta} & \text{Fun}(f \downarrow j, \mathcal{D})
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\text{Res}_f} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
 \text{Res}_j \downarrow & \rho_j \swarrow & \downarrow \text{Res}_{\text{pr}} \\
 \mathcal{C} & \xrightarrow{\Delta} & \text{Fun}(f \downarrow j, \mathcal{C}) \\
 F \downarrow & \text{id}_2 \swarrow & \downarrow F \circ \\
 \mathcal{D} & \xrightarrow{\Delta} & \text{Fun}(f \downarrow j, \mathcal{D})
 \end{array}$$

(where ρ_j is as in Lemma C.5) and note that their pastes agree. We want to show that $(\text{id}_1)_!: \text{Lan}_f \circ (F \circ) \rightarrow (F \circ) \circ \text{Lan}_f$ is an equivalence. For this it is enough to show that $\text{Res}_j \circ (\text{id}_1)_!$ is an equivalence for all $j \in \mathcal{J}$. This is one of the transformations that is composed in the paste $(\text{id}_1)_! * (\rho_j)_!$, which is homotopic to $(\text{id}_2)_! * (\rho_j)_!$ by the pasting law for mates. Since $(\rho_j)_!$ is an equivalence by Lemma C.5 and $(\text{id}_2)_!$ is one by assumption, this finishes the proof. \square

Lemma B.6. *Let \mathcal{I} be a category and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor between ∞ -categories that admit colimits indexed by \mathcal{I} . Then the following conditions are equivalent:*

- a) F preserves left Kan extension along the inclusion $\text{inc}: \mathcal{I} \rightarrow \mathcal{I}^\triangleright$.
- b) F preserves colimits indexed by \mathcal{I} .
- c) F sends $\mathcal{I}^\triangleright$ -indexed colimit diagrams to colimit diagrams.

Proof. By Lemma C.16, if the functor F preserves colimits indexed by \mathcal{I} , then it also preserves left Kan extension along inc . The proof of the same lemma also shows that if F preserves left Kan extension along inc , then it preserves the colimits of all diagrams $\mathcal{I} \rightarrow \mathcal{C}$ that lie in the essential image of $\text{Res}_{\text{inc}}: \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$. But, as inc is fully faithful, we have $\text{Res}_{\text{inc}} \text{Lan}_{\text{inc}} \simeq \text{id}$ and thus all diagrams lie in the essential image of Res_{inc} . This shows the equivalence of the first two conditions.

Now note that, by definition, the functor F preserves left Kan extension along inc if and only if the natural transformation

$$\begin{array}{ccc} \text{Lan}_{\text{inc}} \circ (F \circ) & \xrightarrow{\eta} & \text{Lan}_{\text{inc}} \circ (F \circ) \circ \text{Res}_{\text{inc}} \circ \text{Lan}_{\text{inc}} \\ & & \parallel \\ & & \text{Lan}_{\text{inc}} \circ \text{Res}_{\text{inc}} \circ (F \circ) \circ \text{Lan}_{\text{inc}} \xrightarrow{\varepsilon} (F \circ) \circ \text{Lan}_{\text{inc}} \end{array}$$

is an equivalence, where η and ε are the unit respectively counit of the adjunction $\text{Lan}_{\text{inc}} \dashv \text{Res}_{\text{inc}}$. Since η is an equivalence (as inc is fully faithful), this is equivalent to ε being an equivalence on any diagram in the essential image of $(F \circ) \circ \text{Lan}_{\text{inc}}$. By Lemmas C.1 and B.3, this is equivalent to F sending $\mathcal{I}^\triangleright$ -indexed colimit diagrams to colimit diagrams. \square

Lemma C.17. *Let I, J , and K be simplicial sets, $f: I \rightarrow J$ a map, and \mathcal{C} a weakly left f -extensible ∞ -category that admits colimits indexed by K . Then the functor $\text{Lan}_f: \text{Fun}(I, \mathcal{C}) \rightarrow \text{Fun}(J, \mathcal{C})$ preserves colimits indexed by K .*

Proof. By [LurHTT, Proposition 5.2.3.5] left adjoints preserve colimits. Noting that Lan_f is left adjoint, this implies the statement. \square

Lemma C.18. *Let \mathcal{I} and \mathcal{J} be two categories and \mathcal{C} an ∞ -category that admits colimits indexed by \mathcal{I} and limits indexed by \mathcal{J} . Then $\text{colim}_{\mathcal{I}}: \text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \mathcal{C}$ preserves limits indexed by \mathcal{J} if and only if $\text{lim}_{\mathcal{J}}: \text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$ preserves colimits indexed by \mathcal{I} .*

Proof. We show that, if $\text{colim}_{\mathcal{I}}: \text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \mathcal{C}$ preserves limits indexed by \mathcal{J} , then $\text{lim}_{\mathcal{J}}: \text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$ preserves colimits indexed by \mathcal{I} . The other direction follows dually.

By Lemmas B.6, C.7, and C.14, our assumption implies that $\text{Lan}_{\text{inc}_{\mathcal{I}}}$ preserves right Kan extension along $\text{inc}_{\mathcal{J}}$. Hence there is an equivalence

$$(\text{Lan}_{\text{inc}_{\mathcal{I}}} \circ) \circ \text{Ran}_{\text{inc}_{\mathcal{J}}} \simeq \text{Ran}_{\text{inc}_{\mathcal{J}}} \circ (\text{Lan}_{\text{inc}_{\mathcal{I}}} \circ)$$

of functors $\text{Fun}(\mathcal{J}, \text{Fun}(\mathcal{I}, \mathcal{C})) \rightarrow \text{Fun}(\mathcal{J}^\triangleleft, \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C}))$. This transforms, through a few applications of Lemma C.2, to an equivalence

$$\text{Lan}_{\text{inc}_{\mathcal{I}}} \circ (\text{Ran}_{\text{inc}_{\mathcal{J}}} \circ) \simeq (\text{Ran}_{\text{inc}_{\mathcal{J}}} \circ) \circ \text{Lan}_{\text{inc}_{\mathcal{I}}}$$

of functors $\text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{J}, \mathcal{C})) \longrightarrow \text{Fun}(\mathcal{I}^\triangleright, \text{Fun}(\mathcal{J}^\triangleleft, \mathcal{C}))$. This becomes, after postcomposing with $(\text{Res}_{\blacktriangleleft \mathcal{J}} \circ)$, an equivalence

$$\text{Lan}_{\text{inc}_{\mathcal{I}}} \circ (\lim_{\mathcal{J}} \circ) \simeq (\lim_{\mathcal{J}} \circ) \circ \text{Lan}_{\text{inc}_{\mathcal{I}}}$$

of functors $\text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{J}, \mathcal{C})) \longrightarrow \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C})$ (using Lemma C.7 and that restrictions preserve left Kan extensions). Thus $\lim_{\mathcal{J}}$ sends colimit diagrams indexed by $\mathcal{I}^\triangleright$ to colimit diagrams, which we wanted to show. \square

Lemma C.19. *Let \mathcal{I} be a category, i an object of \mathcal{I} , \mathcal{C} and \mathcal{D} two ∞ -categories that admit colimits indexed by \mathcal{I} , $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor that preserves colimits indexed by \mathcal{I} , and $D: \mathcal{I} \rightarrow \mathcal{C}$ a diagram. Then F applied to the structure map $D(i) \rightarrow \text{colim}_{\mathcal{I}} D$ is an equivalence if and only if the structure map $(F \circ D)(i) \rightarrow \text{colim}_{\mathcal{I}} (F \circ D)$ is.*

Proof. This follows from Lemma B.6 and Remark 1.24. \square

Lemma C.20. *Let $f: I \rightarrow J$ be a map of simplicial sets and $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor between weakly left f -extensible ∞ -categories. Then the diagram*

$$\begin{array}{ccc} (F \circ) & \xrightarrow{\eta^{\mathcal{D}}} & \text{Res}_f \circ \text{Lan}_f \circ (F \circ) \\ \eta^{\mathcal{C}} \downarrow & & \downarrow \chi \\ (F \circ) \circ \text{Res}_f \circ \text{Lan}_f & \xlongequal{\quad} & \text{Res}_f \circ (F \circ) \circ \text{Lan}_f \end{array}$$

commutes up to homotopy, where χ is as in Notation 1.31, and $\eta^{\mathcal{C}}$ and $\eta^{\mathcal{D}}$ are the units of the adjunctions $\text{Lan}_f \dashv \text{Res}_f$ with the respective target categories.

Proof. This is a special case of Lemma A.2. \square

Lemma C.21. *Let \mathcal{I} be a category, $F: \mathcal{C} \rightarrow \mathcal{D}$ a functor between ∞ -categories that admit limits indexed by \mathcal{I} , and denote by $\text{inc}: \mathcal{I} \rightarrow \mathcal{I}^\triangleleft$ the inclusion. Then the following diagram in $\text{Fun}(\text{Fun}(\mathcal{I}^\triangleleft, \mathcal{C}), \mathcal{D})$ commutes up to homotopy*

$$\begin{array}{ccc} F \circ \text{Res}_{\blacktriangleleft} & \longrightarrow & F \circ \lim_{\mathcal{I}} \circ \text{Res}_{\text{inc}} \xrightarrow{\chi} \lim_{\mathcal{I}} \circ (F \circ) \circ \text{Res}_{\text{inc}} \\ \parallel & & \parallel \\ \text{Res}_{\blacktriangleleft} \circ (F \circ) & \longrightarrow & \lim_{\mathcal{I}} \circ \text{Res}_{\text{inc}} \circ (F \circ) \end{array}$$

where the upper left and the bottom horizontal morphism are given by the respective canonical map to the limit, and χ is as in Notation 1.31.

Proof. Consider the two diagrams

$$\begin{array}{ccc}
 \text{Fun}(\mathcal{I}^\triangleleft, \mathcal{C}) & \xrightarrow{\text{id}} & \text{Fun}(\mathcal{I}^\triangleleft, \mathcal{C}) \\
 \text{Res} \downarrow & \nearrow \xi & \downarrow \text{Res}_{\text{inc}} \\
 \mathcal{C} & \xrightarrow{\Delta} & \text{Fun}(\mathcal{I}, \mathcal{C}) \\
 F \downarrow & \nearrow \text{id}_1 & \downarrow F_\circ \\
 \mathcal{D} & \xrightarrow{\Delta} & \text{Fun}(\mathcal{I}, \mathcal{D})
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{Fun}(\mathcal{I}^\triangleright, \mathcal{C}) & \xrightarrow{\text{id}} & \text{Fun}(\mathcal{I}^\triangleleft, \mathcal{C}) \\
 F_\circ \downarrow & \nearrow \text{id}_2 & \downarrow F_\circ \\
 \text{Fun}(\mathcal{I}^\triangleleft, \mathcal{D}) & \xrightarrow{\text{id}} & \text{Fun}(\mathcal{I}^\triangleleft, \mathcal{D}) \\
 \text{Res} \downarrow & \nearrow \xi & \downarrow \text{Res}_{\text{inc}} \\
 \mathcal{D} & \xrightarrow{\Delta} & \text{Fun}(\mathcal{I}, \mathcal{D})
 \end{array}$$

where ξ is as in Notation 1.25, and note that their pastes agree. Now the pasting law for mates implies the desired statement since the mates of the transformations labeled ξ are the canonical maps to the limit, the mate of id_1 is χ , and the mate of id_2 is the identity. \square

D. Generalities

D.1. about posets

Lemma D.1. *Let \mathcal{I} be a poset, \mathcal{C} a category, and $f: \mathcal{I} \rightarrow \mathcal{C}$ a full functor. Then f is injective.*

Proof. Assume that there exist $i \neq i' \in \mathcal{I}$ such that $f(i) = f(i')$. Since f is full there must be both a map $i \rightarrow i'$ and a map $i' \rightarrow i$ being mapped to $\text{id}_{f(i)}$ by f . This contradicts the definition of a poset. \square

Lemma D.2. *Let $f: \mathcal{I} \rightarrow \mathcal{J}$ and $g: \mathcal{J} \rightarrow \mathcal{K}$ be functors between categories such that \mathcal{J} is a poset and $g \circ f$ is full. Then f is full.*

Proof. This follows from the fact that, when a surjective map of sets factors over a set with at most one element, the first map in this factorization is also surjective. \square

Lemma D.3. *Let \mathcal{I} and \mathcal{J} be posets. Assume that both \mathcal{I} and \mathcal{J} have initial objects $\emptyset_{\mathcal{I}}$ respectively $\emptyset_{\mathcal{J}}$ and that \mathcal{I} has a terminal object $*_{\mathcal{I}} \neq \emptyset_{\mathcal{I}}$. Then $(\mathcal{I} \times \mathcal{J})_{>\emptyset}$ is contractible.*

Proof. Since \mathcal{J} has an initial object and is thus contractible, it is enough to show that there is an adjoint pair of functors between $(\mathcal{I} \times \mathcal{J})_{>\emptyset}$ and \mathcal{J} as this implies that they are homotopy equivalent.

To this end, let $l: (\mathcal{I} \times \mathcal{J})_{>\emptyset} \rightarrow \mathcal{J}$ be given by the projection, i.e. $l(i, j) = j$, and $r: \mathcal{J} \rightarrow (\mathcal{I} \times \mathcal{J})_{>\emptyset}$ by $r(j) = (*_{\mathcal{I}}, j)$. Note that r is well-defined as, by assumption, we have $*_{\mathcal{I}} \neq \emptyset_{\mathcal{I}}$. To check that l is indeed left adjoint to r , we need to prove that, for all $(i, j) \in (\mathcal{I} \times \mathcal{J})_{>\emptyset}$ and $j' \in \mathcal{J}$, we have $j = l(i, j) \leq j'$ if and only if $(i, j) \leq r(j') = (*_{\mathcal{I}}, j')$. This is true by the assumption of $*_{\mathcal{I}}$ being terminal in \mathcal{I} . \square

Lemma D.4. *Let $f: \mathcal{S} \rightarrow \mathcal{T}$ and $f': \mathcal{S}' \rightarrow \mathcal{T}$ be maps of posets where $\mathcal{S}, \mathcal{S}' \in \mathbf{Pos}_{\emptyset}$ and $\mathcal{T} \in \mathbf{Pos}_{\square}$. Assume that $f^{-1}(\emptyset_{\mathcal{T}}) = \{\emptyset_{\mathcal{S}}\}$ and $(f')^{-1}(\emptyset_{\mathcal{T}}) = \{\emptyset_{\mathcal{S}'}\}$ and that for all $t \in \mathcal{T}_{>\emptyset}$ one of the posets $f \downarrow t$ and $f' \downarrow t$ has a terminal object which is different from the initial object (in particular this is fulfilled if $f = \text{id}_{\mathcal{T}}$). Then $p: (\mathcal{S} \times \mathcal{S}')_{>\emptyset} \rightarrow \mathcal{T}_{>\emptyset}$ given by $(s, s') \mapsto f(s) \amalg f'(s')$ is homotopy initial.*

Proof. We need that, for all $t \in \mathcal{T}_{>\emptyset}$, the category $p \downarrow t$ is contractible. This comma category can be identified with the full subposet

$$\{(s, s') \in (\mathcal{S} \times \mathcal{S}')_{>\emptyset} \mid f(s) \leq t \text{ and } f'(s') \leq t\} \subseteq (\mathcal{S} \times \mathcal{S}')_{>\emptyset}$$

using the universal property of the coproduct. This, in turn, is isomorphic to the category $((f \downarrow t) \times (f' \downarrow t))_{>\emptyset}$ which is contractible by Lemma D.3. Here, we use that, by our assumptions both $f \downarrow t$ and $f' \downarrow t$ have an initial object ($\emptyset_{\mathcal{S}}$ respectively $\emptyset_{\mathcal{S}'}$) and one of them has a terminal object different from the initial object. \square

D.2. about ∞ -categories

Lemma D.5. *Let $l: \mathcal{C} \rightarrow \mathcal{D}$ and $r: \mathcal{D} \rightarrow \mathcal{C}$ be two functors between ∞ -categories such that l is left adjoint to r with unit η and counit ε . Then, for any simplicial set K , the functor $(l \circ): \text{Fun}(K, \mathcal{C}) \rightarrow \text{Fun}(K, \mathcal{D})$ is left adjoint to $(r \circ): \text{Fun}(K, \mathcal{D}) \rightarrow \text{Fun}(K, \mathcal{C})$ with unit $(\eta \circ)$ and counit $(\varepsilon \circ)$.*

Proof. This is [RV, Proposition 2.1.7 (iii)]. \square

Lemma D.6. *Let the following be a pullback square in the 1-category of simplicial sets*

$$\begin{array}{ccc} K & \longrightarrow & \mathcal{D} \\ f \downarrow & & \downarrow g \\ \mathcal{E} & \longrightarrow & \mathcal{C} \end{array}$$

where \mathcal{D} and \mathcal{E} are ∞ -categories and \mathcal{C} is a category. Then K is an ∞ -category.

Proof. By [LurHTT, Proposition 2.3.1.5], the functor g is an inner fibration. But then f is also an inner fibration since they are stable under pullbacks. Since \mathcal{E} is an ∞ -category, the constant map $c: \mathcal{E} \rightarrow *$ is also an inner fibration. Since inner fibrations are closed under composition, the constant map $K \rightarrow *$ is thus also an inner fibration and hence K an ∞ -category. \square

Lemma D.7. *Let \mathcal{C} be an ∞ -category, \mathcal{D} a category, and $f: \mathcal{D} \rightarrow \text{h}\mathcal{C}$ a functor. Furthermore, let \mathcal{E} be a pullback (in the 1-category of simplicial sets) as in the following diagram*

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{g} & \mathcal{C} \\ p \downarrow & & \downarrow \pi_{\mathcal{C}} \\ \mathcal{D} & \xrightarrow{f} & \text{h}\mathcal{C} \end{array}$$

where $\pi_{\mathcal{C}}$ denotes the canonical functor to the homotopy category. Then \mathcal{E} is an ∞ -category, there is a unique isomorphism $\mathcal{D} \cong \text{h}\mathcal{E}$ under \mathcal{E} , and, for two objects $E, E' \in \mathcal{E}$ and morphism $d: p(E) \rightarrow p(E')$ in \mathcal{D} , the functor g induces an equivalence from the path component of $\text{Map}_{\mathcal{E}}(E, E')$ over d (there is only one such component by the identification $\mathcal{D} \cong \text{h}\mathcal{E}$) to the path component of $\text{Map}_{\mathcal{C}}(g(E), g(E'))$ over $f(d)$.

Proof. That \mathcal{E} is an ∞ -category follows directly from Lemma D.6. Furthermore note that, since $\pi_{\mathcal{C}}$ is a bijection on objects, the map p is as well, i.e. we can identify objects of \mathcal{E} with objects of \mathcal{D} . Now, by the universal property of the pullback the functor g induces, for any morphism $d: D \rightarrow D'$ in \mathcal{D} , an isomorphism from the simplicial subset of $\text{Hom}_{\mathcal{E}}^{\mathbb{R}}(D, D')$ lying over d to the simplicial subset of $\text{Hom}_{\mathcal{C}}^{\mathbb{R}}(f(D), f(D'))$ lying over $f(d)$ (cf. [LurHTT, Section 1.2.2] for the definition of $\text{Hom}^{\mathbb{R}}$). This shows the last statement. To obtain the identification $\mathcal{D} \cong \text{h}\mathcal{E}$, note that what we have already shown implies that the part of $\text{Map}_{\mathcal{E}}(D, D')$ lying over d is path-connected and that these parts are, for different morphisms in \mathcal{D} , disjoint path-components that cover the whole space. \square

Lemma D.8. *Let \mathcal{C} be an ∞ -category. Then the canonical map $\pi_{\mathcal{C}}: \mathcal{C} \rightarrow \mathbf{h}\mathcal{C}$ to its homotopy category is a categorical fibration.*

Proof. By [LurHTT, Corollary 2.4.6.5] the statement is equivalent to $\pi_{\mathcal{C}}$ being an inner fibration such that for every equivalence $f: D \rightarrow D'$ in $\mathbf{h}\mathcal{C}$ and $C \in \mathcal{C}$ with $\pi_{\mathcal{C}}(C) = D$ there exists an equivalence $g: C \rightarrow C'$ in \mathcal{C} such that $\pi_{\mathcal{C}}(g) = f$. That it is an inner fibration follows directly from [LurHTT, Proposition 2.3.1.5] and the other property is clear from the definition of the homotopy category. \square

Lemma D.9. *Denote by S the simplicial set obtained from the directed graph with vertices \mathbb{N}_0 and an edge $n \rightarrow n+1$ for every $n \in \mathbb{N}_0$, and by $i: S \rightarrow \mathbb{N}_0$ the canonical inclusion of simplicial sets. Then, for every ∞ -category \mathcal{C} , the restriction $\text{Res}_i: \text{Fun}(\mathbb{N}_0, \mathcal{C}) \rightarrow \text{Fun}(S, \mathcal{C})$ is a trivial Kan fibration.*

In particular, for every functor $f: S \rightarrow \mathcal{C}$, there is an essentially unique functor $g: \mathbb{N}_0 \rightarrow \mathcal{C}$ such that $g \circ i = f$. Less formally: to specify a sequential diagram in \mathcal{C} it is enough to specify a sequence of composable morphisms $(D_n \rightarrow D_{n+1})_{n \in \mathbb{N}_0}$ in \mathcal{C} .

Proof. This is [LurK, Theorem 00J4] applied to the directed graph used to define S . \square

Lemma D.10. *Let $L: \mathcal{C} \rightarrow \mathcal{D}$ and $R: \mathcal{D} \rightarrow \mathcal{C}$ be two functors between ∞ -categories such that there is an adjunction $L \dashv R$ with unit $\eta: \text{id}_{\mathcal{C}} \rightarrow R \circ L$. Then, for all $c \in \mathcal{C}$ and $d \in \mathcal{D}$, the composition*

$$\text{Map}_{\mathcal{D}}(L(c), d) \xrightarrow{R} \text{Map}_{\mathcal{C}}(RL(c), R(d)) \xrightarrow{\circ\eta(c)} \text{Map}_{\mathcal{C}}(c, R(d))$$

is an equivalence.

Proof. We claim that $(\varepsilon(d) \circ) \circ L$ is a quasi-inverse, where $\varepsilon: L \circ R \rightarrow \text{id}_{\mathcal{D}}$ is the counit of the above adjunction $L \dashv R$. To see that it is a left inverse consider the following homotopy commutative diagram

$$\begin{array}{ccccc} \text{Map}_{\mathcal{D}}(L(c), d) & & \xrightarrow{\circ\varepsilon(L(c))} & & \\ \downarrow R & \searrow LR & & & \\ \text{Map}_{\mathcal{C}}(RL(c), R(d)) & \xrightarrow{L} & \text{Map}_{\mathcal{D}}(LRL(c), LR(d)) & \xrightarrow{\varepsilon(d) \circ} & \text{Map}_{\mathcal{D}}(LRL(c), d) \\ \downarrow \circ\eta(c) & & \downarrow \circ L(\eta(c)) & & \downarrow \circ L(\eta(c)) \\ \text{Map}_{\mathcal{C}}(c, R(d)) & \xrightarrow{L} & \text{Map}_{\mathcal{D}}(L(c), LR(d)) & \xrightarrow{\varepsilon(d) \circ} & \text{Map}_{\mathcal{D}}(L(c), d) \end{array}$$

and note that the composition along the right side of the diagram is homotopic to the identity by one of the triangle identities. Analogously one can show that it is also a right inverse. \square

D.3. about cartesian diagrams

Lemma D.11. *Let \mathcal{I} and \mathcal{J} be categories that have initial objects, $f: \mathcal{I} \rightarrow \mathcal{J}$ an initial object preserving functor, and \mathcal{C} an ∞ -category that admits limits indexed by both $\mathcal{I}_{>\emptyset}$ and by $\mathcal{J}_{>\emptyset}$. Furthermore, assume that f restricts to a functor $\mathcal{I}_{>\emptyset} \rightarrow \mathcal{J}_{>\emptyset}$ and that this functor is homotopy initial.*

Then a diagram $D: \mathcal{J} \rightarrow \mathcal{C}$ is cartesian if and only if $D \circ f: \mathcal{I} \rightarrow \mathcal{C}$ is cartesian.

Proof. We have, by (the dual of) Lemma C.9, a homotopy commutative diagram

$$\begin{array}{ccc} D(\emptyset_{\mathcal{J}}) & \xlongequal{\quad} & (D \circ f)(\emptyset_{\mathcal{I}}) \\ \downarrow & & \downarrow \\ \lim_{\mathcal{J}_{>\emptyset}} D|_{\mathcal{J}_{>\emptyset}} & \xrightarrow[\cong]{f^*} & \lim_{\mathcal{I}_{>\emptyset}} (D \circ f)|_{\mathcal{I}_{>\emptyset}} \end{array}$$

in which the bottom horizontal map is an equivalence by assumption. Hence, the left vertical map is an equivalence if and only if the right vertical map is an equivalence, which we wanted to show. \square

Lemma D.12. *Let \mathcal{I} and \mathcal{J} be categories such that \mathcal{J} has an initial object, \mathcal{C} an ∞ -category that admits limits indexed by $\mathcal{J}_{>\emptyset}$, and $D: \mathcal{I} \times \mathcal{J} \rightarrow \mathcal{C}$ a functor. Denote by $D_{\mathcal{I}}: \mathcal{I} \rightarrow \text{Fun}(\mathcal{J}, \mathcal{C})$ and $D_{\mathcal{J}}: \mathcal{J} \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$ the curried functors. Furthermore assume that $D_{\mathcal{I}}(i) = \text{Res}_i \circ D_{\mathcal{J}}: \mathcal{J} \rightarrow \mathcal{C}$ is cartesian for all $i \in \mathcal{I}$.*

a) *If \mathcal{C} admits limits indexed by \mathcal{I} , then $D_{\mathcal{J}}$, $\lim_{\mathcal{I}} \circ D_{\mathcal{J}}$, and $\lim_{\mathcal{I}} D_{\mathcal{I}}$ are all cartesian.*

b) *If \mathcal{C} admits colimits indexed by \mathcal{I} and the functor $\text{colim}_{\mathcal{I}}: \text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \mathcal{C}$ preserves limits indexed by $\mathcal{J}_{>\emptyset}$, then $\text{colim}_{\mathcal{I}} \circ D_{\mathcal{J}}$ and $\text{colim}_{\mathcal{I}} D_{\mathcal{I}}$ are cartesian.*

Proof. We first show that $D_{\mathcal{J}}$ is cartesian, i.e. that the canonical map $D_{\mathcal{J}}(\emptyset) \rightarrow (\lim_{\mathcal{J}_{>\emptyset}} \text{Res}_{\mathcal{J}_{>\emptyset}})(D_{\mathcal{J}})$ is an equivalence. For this it is enough that its restriction to i is an equivalence for all $i \in \mathcal{I}$, which follows by assumption and Lemma C.21 since Res_i preserves limits.

Now, if \mathcal{C} admits limits indexed by \mathcal{I} , then $\lim_{\mathcal{I}} \circ D_{\mathcal{J}}$ is also cartesian by again Lemma C.21 since $\lim_{\mathcal{I}}$ preserves limits by Lemma C.17. This also implies that $\lim_{\mathcal{I}} D_{\mathcal{I}}$ is cartesian by Lemma C.3. The statement about colimits can be shown in the same way since we only used that $\lim_{\mathcal{I}}$ preserves limits indexed by $\mathcal{J}_{>\emptyset}$. \square

Lemma D.13. *Let \mathcal{I} and \mathcal{J} be categories with initial objects and \mathcal{C} an ∞ -category that admits limits indexed by $\mathcal{J}_{>\emptyset}$. Furthermore, let $D: \mathcal{I} \times \mathcal{J} \rightarrow \mathcal{C}$ be a diagram such that, for each $i \in \mathcal{I}$, the restriction $D|_{\{i\} \times \mathcal{J}}: \mathcal{J} \rightarrow \mathcal{C}$ is cartesian. Then D is a limit diagram.*

Proof. We consider the inclusions

$$\mathcal{I} \times (\mathcal{J}_{>\emptyset}) \xrightarrow{\iota} (\mathcal{I} \times \mathcal{J})_{>\emptyset} \xrightarrow{\kappa} \mathcal{I} \times \mathcal{J}.$$

By assumption and [LurHTT, 4.3.2.9], the functor D is a right Kan extension of $\text{Res}_{\kappa \circ \iota}$ along $\kappa \circ \iota$ in the sense of [LurHTT, 4.3.2.2]. In the same way we also obtain that $\text{Res}_{\kappa} D$ is a right Kan extension of $\text{Res}_{\kappa \circ \iota} D$ along ι . Then, by [LurHTT, 4.3.2.8], the diagram D is a right Kan extension of $\text{Res}_{\kappa} D$ along κ , i.e. D is a limit diagram. \square

References

- [AC] Gregory Arone and Michael Ching. “Goodwillie Calculus”. In: *Handbook of Homotopy Theory*. Ed. by Haynes Miller. To be published. CRC Press, 2020. arXiv: 1902.00803.
- [Ayo] Joseph Ayoub. “Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I.” In: *Astérisque* 314 (2007), pp. x + 466.
- [DP] B. A. Davey and H. A. Priestley. *Introduction to Lattices and Order*. 2nd ed. Cambridge University Press, 2002. DOI: 10.1017/CB09780511809088.
- [Goo90] Thomas G. Goodwillie. “Calculus I: The first derivative of pseudoisotopy theory”. In: *K-Theory* 4.1 (1990), pp. 1–27. DOI: 10.1007/BF00534191.
- [Goo92] Thomas G. Goodwillie. “Calculus II: Analytic functors”. In: *K-Theory* 5.4 (1992), pp. 295–332. DOI: 10.1007/BF00535644.
- [Goo03] Thomas G. Goodwillie. “Calculus III: Taylor series”. In: *Geometry & Topology* 7 (2003), pp. 645–711. DOI: 10.2140/gt.2003.7.645. arXiv: math/0310481.
- [GPS] Moritz Groth, Kate Ponto, and Michael Shulman. “Mayer-Vietoris sequences in stable derivators”. In: *Homology Homotopy Appl.* 16.1 (2014), pp. 265–294. DOI: 10.4310/HHA.2014.v16.n1.a15. arXiv: 1306.2072.
- [KS] G. M. Kelly and Ross Street. “Review of the elements of 2-categories”. In: *Category Seminar (Proceedings Sydney Category Theory Seminar 1972/1973)*. Ed. by Gregory M. Kelly. Vol. 420. Lecture Notes in Mathematics. Springer-Verlag, 1974, pp. 75–103.
- [Kuh] Nicholas J. Kuhn. “Goodwillie towers and chromatic homotopy: an overview”. In: *Proceedings of the Nishida Fest (Kinosaki 2003)*. Vol. 10. Geometry & Topology Monographs. Geometry & Topology Publications, 2007, pp. 245–279. DOI: 10.2140/gtm.2007.10.245. arXiv: math/0410342.
- [LurHTT] Jacob Lurie. *Higher Topos Theory*. Annals of Mathematics Studies 170. Princeton University Press, 2009. arXiv: math/0608040. URL: <http://www.math.harvard.edu/~lurie/papers/HTT.pdf>.
- [LurHA] Jacob Lurie. *Higher Algebra*. Unpublished Manuscript. Version dated September 18, 2017. URL: <http://www.math.harvard.edu/~lurie/papers/HA.pdf>.
- [LurK] Jacob Lurie. *Kerodon*. An online resource for homotopy-coherent mathematics. URL: <https://kerodon.net> (visited on 2019-09-14).
- [May] J. P. May. *A concise course in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, 1999. URL: <https://www.math.uchicago.edu/~may/CONCISE/ConciseRevised.pdf>.
- [Per] Luís Alexandre Pereira. “A general context for Goodwillie Calculus”. Preprint. 2013. arXiv: 1301.2832.

- [Rez] Charles Rezk. “A streamlined proof of Goodwillie’s n -excisive approximation”. In: *Algebraic & Geometric Topology* 13 (2013), pp. 1049–1051. DOI: 10.2140/agt.2013.13.1049. arXiv: 0812.1324.
- [RV15] Emily Riehl and Dominic Verity. “The 2-category theory of quasi-categories”. In: *Advances in Mathematics* 280 (2015), pp. 549–642. DOI: 10.1016/j.aim.2015.04.021. arXiv: 1306.5144.
- [RV] Emily Riehl and Dominic Verity. *Elements of ∞ -Category Theory*. Unpublished Manuscript. Version dated August 28, 2019. URL: <http://www.math.jhu.edu/~eriehl/elements.pdf>.